arXiv:1211.3028v1 [math.GT] 13 Nov 2012

MORSE THEORY FOR LAGRANGE MULTIPLIERS AND ADIABATIC LIMITS

STEPHEN SCHECTER AND GUANGBO XU

Abstract. Given two Morse functions f, μ on a compact manifold M, we study the Morse homology for the Lagrange multiplier function on $M \times \mathbb{R}$ which sends (x, η) to $f(x) + \eta\mu(x)$. Take a product metric on $M \times \mathbb{R}$, and rescale its $\mathbb{R}$-component by a factor λ^2. We show that generically, for large λ, the Morse-Smale-Witten chain complex is isomorphic to the one for f and the metric restricted to $\mu^{-1}(0)$, with grading shifted by one. On the other hand, let $\lambda \to 0$, we obtain another chain complex, which is geometrically quite different but has the same homology as the singular homology of $\mu^{-1}(0)$ and the isomorphism between them is provided by the homotopy by varying λ. Our proofs contain both the implicit function theorem on Banach manifolds and geometric singular perturbation theory.

Contents

1. Introduction 1
2. Morse homology and Morse-Smale-Witten complex of $(\mathcal{F}, g_\lambda)$ 4
3. Adiabatic limit $\lambda \to \infty$ 9
4. Fast-slow system associated to the $\lambda \to 0$ limit 16
5. Adiabatic limit $\lambda \to 0$ 23
6. Basic Morse theory and the chain complex $\mathcal{C}^0$ 34
Appendix A. Geometric singular perturbation theory 36
Appendix B. Choosing the center manifold 41
References 42

1. Introduction

Let M be a compact manifold. Suppose f and μ are Morse functions on M, and 0 is a regular value of μ. Then we consider the Lagrange multiplier function

$$\begin{array}{rccc} \mathcal{F}: & M \times \mathbb{R} & \to & \mathbb{R}, \\ & (x, \eta) & \mapsto & f(x) + \eta\mu(x). \end{array}$$

The critical point set of $\mathcal{F}$ is

$$\mathrm{Crit}(\mathcal{F}) = \left\{ (x, \eta) \mid \mu(x) = 0, df(x) + \eta d\mu(x) = 0 \right\}, \tag{1.1}$$

Date: November 14, 2012.

1991 *Mathematics Subject Classification.* 57R70, 37D15.

Key words and phrases. Morse homology, adiabatic limits, geometric singular perturbation theory, exchange lemma.

and there is a bijection

$$\begin{array}{ccc} \mathrm{Crit}(\mathcal{F}) & \simeq & \mathrm{Crit}\left(f|_{\mu^{-1}(0)}\right), \\ (x,\eta) & \to & x. \end{array}$$

This is a topic which is taught in college calculus.

A deeper story is the Morse theory of $\mathcal{F}$. Take a Riemannian metric g on M and the standard Euclidean metric e on $\mathbb{R}$. Denote by g_1 the product metric $g \oplus e$ on $M \times \mathbb{R}$. Then the gradient vector field of $\mathcal{F}$ with respect to g_1 is

$$\nabla \mathcal{F}(x,\eta) = (\nabla f + \eta \nabla \mu, \mu(x)). \tag{1.2}$$

The differential equation for the negative gradient flow of $\mathcal{F}$ is

$$x' = -\left(\nabla f(x) + \eta \nabla \mu(x)\right), \tag{1.3}$$

$$\eta' = -\mu(x). \tag{1.4}$$

Let $p_\pm = (x_\pm, \eta_\pm) \in \mathrm{Crit}(\mathcal{F})$, and let $\mathcal{M}(p_-, p_+)$ be the moduli space of orbits of the negative gradient flow that approach $p_\pm$ as $t \to \pm\infty$ respectively. (If the differential equation $y' = h(y)$ has the solution $y(t)$ defined on $-\infty < t < \infty$, the corresponding *orbit* is $\{y \mid y = y(t) \text{ for some } t\}$. A time shift of a solution yields another solution with the same orbit. One can speak of the limit of an orbit as $t \to -\infty$ or $t \to \infty$, since the limit is the same for any solution corresponding to the orbit. In $\mathcal{M}(p_-, p_+)$, each orbit from p_- to p_+ is represented by a single point.) Then for generic choice of data (f, μ, g), this moduli space will be a smooth manifold with dimension

$$\dim \mathcal{M}(p_-, p_+) = \mathrm{index}(p_-) - \mathrm{index}(p_+) - 1. \tag{1.5}$$

When the dimension is zero, we can count the number of elements of the corresponding moduli space and use the counting to define the Morse-Smale-Witten complex.

We can show that if we rescale the metric on the $\mathbb{R}$-part, by $g_\lambda := g \oplus \lambda^{-2} e$, $\lambda \in \mathbb{R}^+$ we get the same homology groups, for generic λ for which the chain complex is defined. The system (1.3)–(1.4) is replaced by

$$x' = -\left(\nabla f(x) + \eta \nabla \mu(x)\right), \tag{1.6}$$

$$\eta' = -\lambda^2 \mu(x). \tag{1.7}$$

We then consider the limits of the complex for $(\mathcal{F}, g_\lambda)$ as λ approaches ∞ and zero. Orbits in the two limits will be completely different geometric objects, but the counting of them gives the same homology groups. Indeed, in the limit $\lambda \to \infty$, orbits will converge to those of the negative gradient flow of the pair $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ on the hypersurface $\mu^{-1}(0) \subset M$. This implies that for generic λ, the homology of the chain complex for $(\mathcal{F}, g_\lambda)$ will be isomorphic to the singular homology of $\mu^{-1}(0)$. Hence if the $\lambda \to 0$ limit gives a well-defined chain complex, its homology gives a new perspective to look at the homology of $\mu^{-1}(0)$.

1.1. Outline.

In Section 2, we review basic facts about the Morse-Smale-Witten complex, and we construct this complex $\mathcal{C}^\lambda$ for the function $\mathcal{F}$ and the metric g_λ. Throughout the paper we work with generic data (f, μ, g).

Let p_-, p_+ be two critical points of $\mathcal{F}$, and consider orbits of (1.6)–(1.7) from p_- to p_+ for different λ. The energy of such an orbit is $\mathcal{F}(p_-) - \mathcal{F}(p_+)$, which is λ-independent.

For large λ, because of the finiteness of energy, one can show that all orbits from p_- to p_+ will be confined *a priori* in a small neighborhood of $\mu^{-1}(0) \times \mathbb{R}$. In fact, those orbits will be close to the orbits of the negative gradient flow of $f|_{\mu^{-1}(0)}$. This gives the correspondence between the Morse-Smale-Witten chain complex for $(\mathcal{F}, g_\lambda)$ for large λ, and the complex for $\left(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)}\right)$. In Section 3 we give two proofs of these facts, one using Fredholm theory and one using geometric singular perturbation theory [9]. In Appendix A we review facts from geometric singular perturbation theory that are used in this paper.

On the other hand, when $\lambda \to 0$, orbits from p_- to p_+ will converge to some "singular" orbits. This part of the story is a typical example of fast-slow systems studied in geometric singular perturbation theory. There is only one slow variable, η. We will show that for $\text{index}\,(p_-, \mathcal{F}) - \text{index}\,(p_+, \mathcal{F}) = 1$, if $\lambda > 0$ is small enough, then near each singular orbit from p_- to p_+, there exists a unique orbit of (1.6)–(1.7) from p_- to p_+. This gives a correspondence between the Morse-Smale-Witten chain complex of $(\mathcal{F}, g_\lambda)$ for λ small, and a complex $\mathcal{C}^0$ defined by counting those singular orbits. Details are in Sections 4 and 5.

In Section 6, we sketch various wall-crossing phenomena of the complex $\mathcal{C}^0$ when adding μ a constant c, to recover some elementary facts about Morse theory.

The method we use for the $\lambda \to 0$ limit is similar to the one used in [2]. However, in [2], the slow manifolds are normally hyperbolic, so one can use the classical Exchange Lemma [9]. In the current paper, on the other hand, one cannot avoid the existence of points on the slow manifold where normal hyperbolicity is lost. To overcome this difficulty, one needs a more general exchange lemma, which is provided in [15].

All chain complexes will be chain complexes of $\mathbb{Z}_2$-modules, hence no orientation issues will be considered. We denote $\mathbb{R}_+ = \{t \in \mathbb{R} \mid t > 0\}$ and $\mathbb{Z}_+ = \{n \in \mathbb{Z} \mid n > 0\}$ always, and we use $\text{index}(p, f)$ to denote the Morse index for p as a critical point of f.

1.2. Motivation.

Adiabatic limits appears in many context in geometry in a similar way as the Morse theory we consider here. The direct motivation of this work comes from symplectic geometry, where one wants to study the adiabatic limits of the symplectic vortex equation. We briefly discuss the analogy here.

Suppose (X, ω) is a closed symplectic manifold, together with a Hamiltonian G-action with moment map $\mu : M \to \mathfrak{g}^*$, where G is a compact Lie group and $\mathfrak{g}$ is its Lie algebra. Let $H_t : X \to \mathbb{R}$ be a time-dependent, G-invariant Hamiltonian, with Hamiltonian vector field Y_{H_t}. Then consider the action functional (appeared in [3]) on the free loop space of $X \times \mathfrak{g}$,

$$\mathcal{A}_{\mu,H}(x, \xi) = -\int_{\mathbb{D}} u^*\omega + \int_0^1 (H_t(x(t)) + \xi_t \mu(x(t)))dt.$$

Following the standard procedure of defining Floer homology groups, we can construct a similar theory for $\mathcal{A}_{\mu,H}$. The critical point set of $\mathcal{A}_{\mu,H}$ is

$$\mathrm{Crit}(\mathcal{A}_{\mu,H}) = \left\{ (x,\eta) : S^1 \to M \times \mathfrak{g} \mid \mu(x(t)) \equiv 0,\ \dot{x}(t) = Y_{H_t}(t) + \mathcal{X}_{\eta(t)}(x(t)),\ t \in S^1 \right\}.$$

The equation for the negative gradient flow for this action functional, using a G-invariant ω-compatible almost-complex structure J, is

$$\left\{ \begin{array}{rcl} \partial_s u + J\left(\partial_t u - Y_{H_t} - \mathcal{X}_{\eta(t)}\right) & = & 0; \\ \partial_s \eta + \lambda^2 \mu \circ u & = & 0. \end{array} \right. \tag{1.8}$$

This is the *symplectic vortex equation* with a Hamiltonian perturbation H_t, on the infinite cylinder which has the area form $\lambda^2 dsdt$. The corresponding theory can be regarded as an infinite-dimensional equivariant version of the Morse theory we consider in this article. Its $\lambda \to \infty$ adiabatic limit covers (in principle) the Floer homology of the symplectic quotient $\mu^{-1}(0)/G$, which is an example of the general expectation that equivariant symplectic invariants defined from the moduli spaces of solutions to the vortex equation recover the corresponding nonequivariant invariants of the symplectic quotient, via the $\lambda \to \infty$ adiabatic limit (see for example [7]).

It should be useful to study the behavior of solutions in the opposite adiabatic limit in this infinite-dimensional setting. For example, in the setting of Hamiltonian Gromov-Witten invariants (or gauged Gromov-Witten invariants), González-Woodward studied this limit in genus zero case (see [8]). The more general case for this limit is also worth studying. For example, if we take $M = \mathbb{C}^{Nk}$ and $G = U(k)$ such that the symplectic quotient gives the complex Grassmannian $G(k,N)$, Witten showed by a physics argument that using the $\lambda \to 0$ limit one can derive an equivalence of the Gromov-Witten invariants of $G(k,N)$ and the Verlinde algebra associated to $U(k)$ (see [18]).

1.3. **Acknowledgements.** The first author's work was partially supported by NSF under award DMS-1211707.

The second author would like first to thank his advisor Professor Gang Tian, for his help and encouragement, guidance and corrections. He also would like to thank Urs Frauenfelder for many helpful discussions, to thank Robert Lipshitz and Dietmar Salamon for their interest in this work, to thank Hongbin Sun for answering topology questions. At the early stage of this project, the second named author was funded by Professor Helmut Hofer for the Extramural Summer Research Project of Princeton Univeristy in Summer 2011 and he would like to express his gratitude for Professor Hofer's generosity.

2. Morse homology and Morse-Smale-Witten complex of $(\mathcal{F}, g_\lambda)$

2.1. **Morse homology.** For the topological aspects of Morse theory, there is the classical book [14]. Here we will adopt the viewpoint of Witten [17], which is by now standard.

Let M be a smooth manifold without boundary. A smooth function $f : M \to \mathbb{R}$ is a *Morse function* if its differential df is a transverse section of the cotangent bundle T^*M. This implies that the critical points of f are discrete, and by the Morse lemma near each

$p \in \mathrm{Crit}(f)$ (the critical point set), there exists a local coordinate chart $(x_1, \ldots, x_n)$ such that

$$f(x) = f(p) - \sum_{i=1}^{k_p} x_i^2 + \sum_{i=k_p+1}^{n} x_i^2.$$

The integer k_p is well-defined and is called the *Morse index* of the critical point p. We denote this as

$$\mathrm{index}\,(p, f) = k_p \in \mathbb{Z}_+,$$

and we denote the set of critical points of f of index k by $\mathrm{Crit}_k(f)$ Now for any complete Riemannian metric g on M, the gradient vector field ∇f is the dual of the 1-form df. We denote by $\phi_t : M \to M$ the flow of $-\nabla f$. For each $p \in \mathrm{Crit}(f)$, the unstable and stable manifolds are defined by

$$W^u(p) = \left\{ x \in M | \lim_{t \to -\infty} \phi_t(x) = p \right\}, \quad W^s(p) = \left\{ x \in M | \lim_{t \to +\infty} \phi_t(x) = p \right\}$$

and

$$\dim W^u(p) = \mathrm{index}\,(p, f),\ \dim W^s(p) = n - \mathrm{index}\,(p, f).$$

The pair (f, g) is called *Morse-Smale* if for any two critical points p, q, $W^u(p)$ and $W^s(q)$ intersect transversely in M. This condition condition holds for generic (f, g) [16]. For $p \neq q$, there is a free $\mathbb{R}$-action on $W^u(p) \cap W^s(q)$ by time translation along the flow.

Then we define the *Morse-Smale-Witten chain complex* of $\mathbb{Z}_2$-modules for a Morse-Smale pair (f, g),

$$\mathcal{C}^M(f, g) = (C_*, \partial)\,,\ C_k = \bigoplus_{p \in \mathrm{Crit}_k(f)} \mathbb{Z}_2 \langle p \rangle, \tag{2.1}$$

where $\partial : C_k \to C_{k-1}$ is given by

$$\partial p = \sum_{q \in \mathrm{Crit}_{k-1}(f)} n_{p,q} \cdot q,\ n_{p,q} = \#\left[(W^u(p) \cap W^s(q))/\mathbb{R} \right] \in \mathbb{Z}_2. \tag{2.2}$$

One can show that $\#\left[(W^u(p) \cap W^s(q))/\mathbb{R} \right]$ is finite and $\partial \circ \partial = 0$ (under the Palais-Smale condition on f, see [16]), hence $\mathcal{C}^M(f, g)$ is a chain complex. Its homology group is called the *Morse homology* of (f, g) with $\mathbb{Z}_2$-coefficients, denoted by $HM_*(f, g; \mathbb{Z}_2)$. In the case where M is compact, this homology is independent of the choice of Morse-Smale pair (f, g) and is actually isomorphic to the singular homology of M.

2.2. **Morse homology and the analytic setting .**

The book [16] give a comprehensive treatment of the analytical perspective on finite-dimensional Morse theory. Here we give a brief review.

Let p and q be two critical points of f, and consider the nonlinear ODE

$$x' = -\nabla f(x), \tag{2.3}$$

with the boundary conditions

$$\lim_{t\to-\infty} x(t) = p, \quad \lim_{t\to+\infty} x(t) = q. \tag{2.4}$$

The space of solutions to this boundary value problem can be identified with $W^u(p) \cap W^s(q)$ by identifying the solution $x(t)$ with the point $x(0)$.

Indeed, the space of solutions to such a boundary value problem can be viewed as a finite-dimensional submanifold of an infinite-dimensional Banach manifold. We consider the Banach manifold $\mathcal{B}$ of those $W^{1,2}_{loc}$ maps $x : \mathbb{R} \to M$ for which there exists $T > 0$ such that for $t \in [T, +\infty)$ (respectively $t \in (-\infty, -T]$), $d(x(t), q)$ (respectively $d(x(t), p)$) is less than the injectivity radius of (M, g) and $\exp_q^{-1}(x(t)) \in W^{1,2}([T, +\infty), T_qM)$ (respectively $\exp_p^{-1}(x(t)) \in W^{1,2}((-\infty, -T], T_pM)$. There is a Banach space bundle $\mathcal{E} \to \mathcal{B}$ whose fibre over $x : \mathbb{R} \to M$ is $L^2(\mathbb{R}, x^*TM)$. There is a smooth section

$$\begin{aligned} \mathcal{S} :\ \ \mathcal{B}\ &\to\ \ \ \ \mathcal{E} \\ x\ &\mapsto\ x' + \nabla f(x) \end{aligned}$$

whose zero locus is exactly the moduli space $\widetilde{\mathcal{M}}(p, q) \simeq W^u(p) \cap W^s(q)$, the space of solutions to (2.3)–(2.4). This is a Fredholm section, which means that the linearization of $\mathcal{S}$ at each $x \in \mathcal{S}^{-1}(0)$, which is given by

$$\begin{aligned} D\mathcal{S}_x :\ \ T_x\mathcal{B}\ &\to\ \ \ \ \mathcal{E}_x \\ V\ &\mapsto\ \nabla_t V + \nabla_V(\nabla f) \end{aligned}$$

is a Fredholm operator with Fredholm index equal to $\operatorname{index}(p) - \operatorname{index}(q)$.

In order ensure that $\widetilde{\mathcal{M}}(p, q)$ is a smooth manifold, we need that the linearization of $\mathcal{S}$ is surjective along $\mathcal{S}^{-1}(0)$. In this case, we say that $\widetilde{\mathcal{M}}(p, q) = \mathcal{S}^{-1}(0)$ is *transverse*. This condition is equivalent to transversality of $W^u(p)$ and $W^s(q)$.

$\widetilde{\mathcal{M}}(p, q)$ has a free $\mathbb{R}$-action by time translation, so we can define $\mathcal{M}(p, q) = \widetilde{\mathcal{M}}(p, q)/\mathbb{R}$, the space of orbits from p to q. The compactness of $\mathcal{M}(p, q)$ is an issue, especially when M is noncompact. A control on the behavior of a solution $x : \mathbb{R} \to M$ is its *energy*

$$E(x) = \int_{\mathbb{R}} |\nabla f(x(t))|^2 \, dt = f(p) - f(q), \tag{2.5}$$

which only depends on the values of f at p and q. The energy is essential for many useful estimates. An assumption used to yield compactness is the Palais-Smale condition on f (which is automatic when M is compact): any sequence $x_i \in M$ for which $f(x_i)$ is uniformly bounded and $|\nabla f(x_i)| \to 0$ has a convergent subsequence. Under the transversality and Palais-Smale assumptions, if $\operatorname{index}(q, f) = \operatorname{index}(p, f) - 1$, then $\mathcal{M}(p, q)$ is finite; if $\operatorname{index}(q, f) = \operatorname{index}(p, f) - 2$, then $\mathcal{M}(p, q)$ can be compactified to be a smooth one-dimensional manifold with boundary; the boundary points correspond to the broken orbits from p to q that pass through another critical point r with $\operatorname{index}(r, f) = \operatorname{index}(p, f) - 1$. These two facts imply that the boundary operator (2.2) is well-defined and $\partial \circ \partial = 0$.

2.3. **The Morse-Smale-Witten complex of $(\mathcal{F}, g_\lambda)$.**

For the pair $(\mathcal{F}, g_\lambda)$ defined in Section 1, some modifications to the construction of the Morse-Smale-Witten complex are required. We shall assume

(A1) M is a compact manifold.
(A2) f and μ are Morse functions on M.
(A3) 0 is a regular value of μ and $f|_{\mu^{-1}(0)}$ is Morse.

The metric g on M and the Morse function μ satisfying (A3) are arbitrary and fixed. We shall make further assumptions, always denoted (Ak), that are true for generic f. Alternatively, we could fix the Morse functions f; the assumptions are then true for generic (μ, g).

In each of our results, we shall assume without comment that all assumptions (Ak) made up to that point hold.

Lemma 2.1. *Under the assumption (A3), the Lagrange multiplier $\mathcal{F}$ is a Morse function on $M \times \mathbb{R}$ and for any $p = (x_p, \eta_p) \in \operatorname{Crit}(\mathcal{F})$, $\operatorname{index}(p, \mathcal{F}) = \operatorname{index}(x_p, f|_{\mu^{-1}(0)}) + 1$.*

Proof. Let $p \in \operatorname{Crit}(\mathcal{F})$. Since 0 is a regular value of μ, there exists a local coordinate chart $(x_1, \dots, x_n)$ near x_p such that $\mu(x_1, \dots, x_n) = x_n$ and p has coordinates $(0, \dots, 0)$. Then we see that the second derivative of $\mathcal{F}$ at p with respect to the coordinates $(x_1, \dots, x_n, \eta)$ is of the form

$$D^2\mathcal{F}(0) = \begin{pmatrix} A & b & 0 \\ b^T & c & 1 \\ 0 & 1 & 0 \end{pmatrix}. \tag{2.6}$$

Here A is equal to the second derivative of $f|_{\mu^{-1}(0)}$ at x_p with respect to the coordinates $(x_1, \dots, x_{n-1})|_{\mu^{-1}(0)}$. So this implies that $D^2\mathcal{F}$ is nondegenerate at p, and its determinant is equal to $-\det A$. □

Now we consider the unstable and stable manifolds of critical points of $\mathcal{F}$. Since we are dealing with the family of metrics g_λ, we want transversality of unstable and stable manifolds not just for a single g_λ, but as much as possible for the family. Let ϕ_t^λ denote the flow of (1.6)–(1.7) for the given value of λ, let $p_-, p_+ \in \operatorname{Crit}(\mathcal{F})$, and let I be an interval in λ-space. Define the sets

$$\begin{aligned} W^u(p_-, \lambda) &= \left\{ p \in M \times \mathbb{R} \mid \lim_{t \to -\infty} \phi_t^\lambda(p) = p_- \right\}, \\ W^s(p_+, \lambda) &= \left\{ p \in (M \times \mathbb{R} \mid \lim_{t \to +\infty} \phi_t^\lambda(p) = p_+ \right\}, \\ W^u(p_-, I) &= \{(p, \lambda) \in (M \times \mathbb{R}) \times I \mid p \in W^u(p_-, \lambda)\}, \\ W^s(p_+, I) &= \{(p, \lambda) \in (M \times \mathbb{R}) \times I \mid p \in W^s(p_+, \lambda)\}; \end{aligned}$$

$$\widetilde{\mathcal{M}}^I(p_-, p_+) = W^u(p_-, I) \cap W^s(p_+, I), \;\; \mathcal{M}^I(p_-, p_+) = \widetilde{\mathcal{M}}^I(p_-, p_+)/\mathbb{R}.$$

We assume:

(A4) For all $p_-, p_+ \in \operatorname{Crit}(\mathcal{F})$, $W^u(p_-, \mathbb{R}_+)$ and $W^s(p_+, \mathbb{R}_+)$ are transverse.

Assumption (A4) implies that for all but discretely many $\lambda \in \mathbb{R}_+$, $W^u(p_-, \lambda)$ and $W^s(p_+, \lambda)$ are transverse. We denote by $\Lambda^{reg} \subset \mathbb{R}_+$ the subset of λ's for which $\widetilde{\mathcal{M}}^\lambda(p_-, p_+)$ is transverse for all $p_-, p_+ \in \operatorname{Crit}(\mathcal{F})$.

Since the manifold $M \times \mathbb{R}$ is noncompact, we also need the following result:

Lemma 2.2. *For each $\lambda \in \mathbb{R}_+$, the function $\mathcal{F}$ satisfies the Palais-Smale condition with respect to the metric g_λ, i.e., for any sequence $\widetilde{x}_i \in M \times \mathbb{R}$ such that $\mathcal{F}(\widetilde{x}_i)$ is bounded and $\left|\nabla \mathcal{F}(\widetilde{x}_i)\right|_{g_\lambda} \to 0$, there exists a convergent subsequence of $\widetilde{x}_i$.*

Proof. Suppose $\widetilde{x}_i = (x_i, \eta_i)$. Since M is compact, we may assume that $\eta_i \to +\infty$ or $-\infty$. The condition that $\mathcal{F}(\widetilde{x}_i)$ is bounded implies that $\mu(x_i) \to 0$; but since 0 is a regular value of μ, $\nabla \mathcal{F}(\widetilde{x}_i) = \nabla f(x_i) + \eta_i \nabla \mu(x_i)$ cannot have arbitrary small norm. □

Now, for any $\lambda \in \Lambda^{reg}$, we can define the associated Morse-Smale-Witten complex of $(\mathcal{F}, g_\lambda)$, which is denoted by $\mathcal{C}^\lambda = \mathcal{C}(\mathcal{F}, g_\lambda)$. Its homology is denoted by H^λ. All $\mathcal{C}^\lambda$ share the same generators and gradings, but the boundary operator ∂^λ may change when λ crosses one of the exceptional values in $\mathbb{R}_+ \setminus \Lambda^{reg}$. To show that the homology H^λ is independent of λ, we have

Lemma 2.3. *For any $L > 0$, there exists $K_L > 0$ such that for any $p_\pm \in \mathrm{Crit}(\mathcal{F})$, $W^u(p_-, (0, L]) \cap W^s(p_+, (0, L]) \subset M \times [-K_L, K_L]$.*

Proof. Suppose this statement doesn't hold, then there exists $p_-, p_+ \in \mathrm{Crit}(\mathcal{F})$, a sequence λ_i converging to $\lambda_\infty \in [0, L]$ and $(x_i, \eta_i) \in W^u(p_-, \lambda_i) \cap W^s(p_+, \lambda_i)$ such that $\lim_{i\to\infty} |\eta_i| = +\infty$. Suppose for each i there are solutions $\widetilde{x}_i = (x_i, \eta_i) : \mathbb{R} \to M \times \mathbb{R}$ to (1.6), (1.7) for $\lambda = \lambda_i$ such that $(x_i, \eta_i) = \widetilde{x}_i(0)$. Then since λ_i are uniformly bounded, for any $R > 0$, we have

$$|\eta_i|\,|_{[-R,R]} \to \infty \tag{2.7}$$

uniformly.

On one hand, we have

$$\mathcal{F}(p_-) \geq \mathcal{F}(x_i, \eta_i) = f(x_i) + \eta_i \mu(x_i) \geq \mathcal{F}(p_+) \tag{2.8}$$

which implies that $\mu(x_i) \to 0$ uniformly on $[-R, R]$. Then, by the definition of the energy of the solution $\widetilde{x}_i$, we see

$$\mathcal{F}(p_+) - \mathcal{F}(p_-) \geq \int_{-R}^{R} \left\| \nabla f(x_i(t)) + \eta_i(t) \nabla \mu(x_i(t)) \right\|^2 dt \to \infty \tag{2.9}$$

which is impossible. □

Corollary 2.4. *For any $\lambda_1, \lambda_2 \in \Lambda^{reg}$, there is a canonical isomorphism $\Phi_{\lambda_1,\lambda_2} : H^{\lambda_1} \to H^{\lambda_2}$ such that for $\lambda_1, \lambda_2, \lambda_3 \in \Lambda^{reg}$, $\Phi_{\lambda_2,\lambda_3} \circ \Phi_{\lambda_1,\lambda_2} = \Phi_{\lambda_1,\lambda_3}$.*

Proof. For any $\lambda_1, \lambda_2 \in \Lambda^{reg}$, $\lambda_1 < \lambda_2$, we can compare the two complexes $\mathcal{C}^{\lambda_1}$ and $\mathcal{C}^{\lambda_2}$ as in the case of compact manifolds, thanks to the compactness provided by the previous lemma. More precisely, we can either use the continuation principle, or the bifurcation analysis as did in [4] to show that the two chain complexes have isomorphic homology. Note that as we vary λ from λ_1 to λ_2, the critical point set is fixed so there is no "birth-death" of critical points. What may happen is for some $\lambda \in (\lambda_1, \lambda_2)$, there may exists a trajectory between two different critical points with the same Morse index. □

Remark 2.5. In general, the Morse homology of a pair (f, g) will be very sensitive to the pair if the underlying manifold is noncompact (see for example [11]).

3. Adiabatic limit $\lambda \to \infty$

We assume

(A5) $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ is Morse-Smale.

Theorem 3.1. *For $\lambda \in \Lambda^{reg}$ sufficiently large, the complex $\mathcal{C}^\lambda$ is isomorphic to the Morse-Smale-Witten complex of the pair $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ with grading shifted by one.*

Corollary 3.2. *For any $\lambda \in \Lambda^{reg}$, there is a canonical isomorphism*

$$\Phi_{\lambda,+\infty} : H_*^\lambda \to H_{*-1}(\mu^{-1}(0), \mathbb{Z}_2).$$

We give two proofs of Theorem 3.1, one using the infinite-dimensional implicit function theorem and one using geometric singular perturbation theory. The first is more likely to generalize. The second gives more geometric intuition.

3.1. **Proof of Theorem 3.1 using the implicit function theorem.**

3.1.1. *Convergence to orbits in the level set.* For any $\widetilde{x} = (x, \eta) \in \widetilde{\mathcal{M}}^\lambda(p_-, p_+)$, we have

$$E = \mathcal{F}(p_-) - \mathcal{F}(p_+) = -\int_{-\infty}^{+\infty} \left\langle \nabla^\lambda \mathcal{F}(\widetilde{x}(t)), \widetilde{x}'(t) \right\rangle_\lambda dt = \|x'\|_{L^2}^2 + \lambda^2 \|\mu(x)\|_{L^2}^2 . \tag{3.1}$$

So

$$\|\mu(x)\|_{L^2} \leq \frac{E^{\frac{1}{2}}}{\lambda}. \tag{3.2}$$

On the other hand, one has

$$\left\| \frac{d}{dt}\mu(x(t)) \right\|_{L^2} \leq \|d\mu\|_{L^\infty} \|x'(t)\|_{L^2} . \tag{3.3}$$

So by Sobolev embedding $W^{1,2} \to C^0$, for any $\epsilon > 0$, there exists $\Lambda_\epsilon > 0$ such that for $\lambda > \Lambda_\epsilon$ and $\widetilde{x} \in \widetilde{\mathcal{M}}^\lambda(p_-, p_+)$, we have $x(t) \in U_\epsilon = \mu^{-1}((-\epsilon, \epsilon))$.

Now, consider the Banach manifold $\mathcal{B} = \mathcal{B}^{1,2}$ of $W^{1,2}_{loc}$-maps $\widetilde{x} = (x, \eta)$ from $\mathbb{R} \to U_\epsilon \times \mathbb{R}$ such that $\widetilde{x}$ is assymptotic to $(p_\pm, \eta_{p_\pm}) \in \mathrm{Crit}(\mathcal{F})$ at $\pm\infty$ in the following sense: there exists $R > 0$ and $\widetilde{W}_- \in W^{1,2}((-\infty, -R], T_{p_-}M \oplus \mathbb{R})$ and $\widetilde{W}_+ \in W^{1,2}([R, +\infty), T_{p_+}M \oplus \mathbb{R})$ such that $\widetilde{x}|_{(-\infty,-R]} = \exp_{p_-} \widetilde{W}_-$, $\widetilde{x}|_{[R,+\infty)} = \exp_{p_+} \widetilde{W}_+$. Consider $\mathcal{E} \to \mathcal{B}$ the Banach space bundle whose fibre over $\widetilde{x}$ is $\mathcal{E}_{\widetilde{x}} = L^2(x^*TM \oplus \mathbb{R})$. For $\lambda > \Lambda_\epsilon$, consider the Fredholm section $\widetilde{S}^\lambda : \mathcal{B} \to \mathcal{E}$ given by

$$\widetilde{S}^\lambda(x, \eta) = \left(\frac{dx}{dt} + \nabla f + \eta \nabla\mu, \frac{d\eta}{dt} + \lambda^2 \mu(x) \right). \tag{3.4}$$

Then $\widetilde{\mathcal{M}}^\lambda(p_-, p_+) = \left(\widetilde{S}^\lambda\right)^{-1}(0)$. The linearization of $\widetilde{\mathcal{S}}^\lambda$ at $\widetilde{x} = (x, \eta) \in \mathcal{B}$ is given by

$$\begin{array}{cccc} \widetilde{\mathcal{D}}_{\widetilde{x}} : & T_{\widetilde{x}}\mathcal{B} & \to & \mathcal{E}_{\widetilde{x}} \\ & \begin{pmatrix} V \\ H \end{pmatrix} & \mapsto & \begin{pmatrix} \nabla_t V + \nabla_V(\nabla f + \eta\nabla\mu) + H\nabla\mu \\ \frac{dH}{dt} + \lambda^2 d\mu(V) \end{pmatrix} \end{array} \tag{3.5}$$

On U_ϵ, there is the line bundle $L \subset TM$ generated by $\nabla\mu$, and denote by $L^\perp$ the orthogonal complement with respect to the Riemannian metric g. Then, for any $\widetilde{x}$, we can decompose the domain and target space of $\widetilde{D}_{\widetilde{x}}$ as

$$T_{\widetilde{x}}\mathcal{B} \simeq W_L(\widetilde{x}) \oplus W_T(\widetilde{x}),\ W_L(\widetilde{x}) = W^{1,2}(x^*L \oplus \mathbb{R}),\ W_T(\widetilde{x}) = W^{1,2}(x^*L^\perp); \tag{3.6}$$

$$\mathcal{E}_{\widetilde{x}} \simeq \mathcal{E}_L(\widetilde{x}) \oplus \mathcal{E}_T(\widetilde{x}),\ \mathcal{E}_L(\widetilde{x}) = L^2(x^*L \oplus \mathbb{R}),\ \mathcal{E}_T(\widetilde{x}) = L^2(x^*L^\perp). \tag{3.7}$$

We rescale the norms on $W_L(\widetilde{x})$ and $\mathcal{E}_L(\widetilde{x})$ as follows. We identify $(h_1\nabla\mu, h_2) \in W_L(\widetilde{x})$ with $(h_1, h_2) \in W^{1,2} \oplus W^{1,2}$ and define

$$\|(h_1, h_2)\|_{W_\lambda} = \lambda \|h_1\|_{L_2} + \|h_1'\|_{L^2} + \|h_2\|_{L^2} + \lambda^{-1}\|h_2'\|_{L^2}; \tag{3.8}$$

and for $(h_1\nabla\mu, h_2) \in \mathcal{E}_L(\widetilde{x})$, identify it with $(h_1, h_2) \in L^2 \oplus L^2$ and define

$$\|(h_1, h_2)\|_{L_\lambda} = \|h_1\|_{L^2} + \lambda^{-1}\|h_2\|_{L^2}. \tag{3.9}$$

We leave the norms on their complements unchanged, and use W_λ and L_λ to denote the norms on $T_{\widetilde{x}}\mathcal{B}$ and $\mathcal{E}_{\widetilde{x}}$ respectively.

Now we describe the limit objects. There exists a smooth function $\widetilde{\eta} : U_\epsilon \to \mathbb{R}$ defined by the condition

$$\langle \nabla\mu(x), \nabla f(x) + \widetilde{\eta}(x)\nabla\mu(x)\rangle = 0. \tag{3.10}$$

Then $\nabla f + \widetilde{\eta}\nabla\mu$ is a smooth vector field whose restriction to $\mu^{-1}(0)$ is the gradient of $\overline{f} = f|_{\mu^{-1}(0)}$ with respect to the restriction of the Riemannian metric. We denote by $\widetilde{\mathcal{N}}^\infty(p_-, p_+)$ the space of solutions of the negative gradient flow of $\overline{f}$, and by $\mathcal{N}^\infty(p_-, p_+)$ the quotient space by identifying reparametrizations, and by $\overline{\mathcal{N}}^\infty(p_-, p_+)$ the compactified moduli space by adding broken orbits.

Proposition 3.3. *Suppose $\lambda_\nu \to \infty$ and $\widetilde{x}_\nu = (x_\nu, \xi_\nu) \in \widetilde{\mathcal{M}}^{\lambda_\nu}(p_-, p_+)$. Then there is a subsequence, still indexed by ν, and a broken trajectory $\mathcal{Y} = ([y_i])_{i=1}^n \in \overline{\mathcal{N}}^\infty(p_-, p_+)$ and a constant $c_0 > 0$ such that*

1. $\|(\mu(x_\nu), \eta_\nu - \widetilde{\eta}(x_\nu))\|_{W_\lambda} \leq c_0\lambda_\nu^{-1}$;
2. *There exists $t_{1,\nu}, t_{2,\nu}, \dots, t_{n,\nu} \in \mathbb{R}$ such that $x_\nu(t_{i,\nu} + \cdot)$ converges to y_i in C^1_{loc}-topology;*

Proof. Apply $d\mu$ to the equation $x_\nu'(t) + \nabla f(x_\nu(t)) + \eta_\nu(t)\nabla\mu(x_\nu(t)) = 0$, we obtain

$$\begin{aligned} 0 = \frac{d}{dt}\mu(x_\nu(t)) &+ \langle \nabla\mu(x_\nu(t)), \nabla f(x_\nu(t)) + \eta_\nu(t)\nabla\mu(x_\nu(t))\rangle \\ &= \frac{d}{dt}\mu(x_\nu(t)) + \lambda_\nu |\nabla\mu|^2 \left(\frac{1}{\lambda_\nu}(\eta_\nu(t) - \widetilde{\eta}(x_\nu(t)))\right). \end{aligned} \tag{3.11}$$

Also we have

$$\frac{d}{dt}\left(\frac{1}{\lambda_\nu}(\eta_\nu(t) - \widetilde{\eta}(x_\nu(t)))\right) + \lambda\mu(x_\nu(t)) = -\frac{1}{\lambda_\nu}\frac{d}{dt}\widetilde{\eta}(x_\nu(t)). \tag{3.12}$$

Consider the linear operator

$$\begin{array}{rccc} D: & W^{1,2}(\mathbb{R}, \mathbb{R}^2) & \to & L^2(\mathbb{R}, \mathbb{R}^2) \\ & (f_1, f_2) & \mapsto & \left(\frac{d}{dt}f_1 + \lambda|\nabla\mu|^2 f_2, \frac{d}{dt}f_2 + \lambda f_1\right) \end{array}. \tag{3.13}$$

If regarded as an unbounded operator from L^2 to L^2, then it is bounded from below by $c\lambda$ for some constant c. Then (3.11) and (3.12) imply that

$$\|\mu(x_\nu)\|_{L^2} + \lambda_\nu^{-1} \|\eta_\nu - \widetilde{\eta}(x_\nu)\|_{L^2} \leq \frac{c}{\lambda_\nu^2} \left\| \frac{d}{dt} \widetilde{\eta}(x_\nu) \right\|_{L^2} \leq c\lambda_\nu^{-2}. \tag{3.14}$$

Also, D has a uniformly bounded right inverse, so

$$\|\mu(x_\nu)\|_{W^{1,2}} + \lambda_\nu^{-1} \|\eta_\nu - \widetilde{\eta}(x_\nu)\|_{W^{1,2}} \leq c\lambda_\nu^{-1}. \tag{3.15}$$

This implies the first claim of this proposition. Moreover, the Sobolev embedding $W^{1,2} \to C^0$ implies in particular that η_ν is uniformly bounded since $\widetilde{\eta}(x_\nu)$ is. It follows that $|x_\nu'|$ is uniformly bounded.

Now we identify $U_\epsilon \simeq \mu^{-1}(0) \times (-\epsilon, \epsilon)$ such that the projection to the second component is equal to μ. Then we can write

$$x_\nu(t) = (\overline{x}_\nu(t), \mu(x_\nu(t))). \tag{3.16}$$

Projecting to the first factor and using the fact that $|x_\nu'|$ is uniformly bounded, we see that there exists $K > 0$ independent of ν, such that

$$\left| \overline{x}_\nu'(t) + \nabla \overline{f} \left((\overline{x}_\nu(t)) \right) \right| \leq K |\mu(x_\nu(t))|. \tag{3.17}$$

This implies that a subsequence of $\overline{x}_\nu$ converges to a broken trajectory for the induced function $\overline{f}$ in C^0_{loc}-topology. More precisely, there exists $t_{1,\nu}, \ldots, t_{k,\nu} \in \mathbb{R}$ such that $\overline{x}_\nu(t_{i,\nu} + \cdot)$ converges to a trajectory y_i in $\mu^{-1}(0)$ in C^0_{loc}-topology.

Then, by the Sobolev embedding $W^{1,2} \to C^0$ and the C^0_{loc} convergence of $x_\nu(t_{i,\nu} + \cdot)$ to y_i, we see that $\eta_\nu(t_{i,\nu} + \cdot)$ converges to $\widetilde{\eta}(y_i)$ in C^0_{loc}. This implis that the convergence $x_\nu(t_{i,\nu} + \cdot)$ to y_i is in C^1_{loc}. □

3.1.2. *Applying implicit function theorem and the isomorphism of chain complexes.* Now we want to prove that, any $y \in \widetilde{\mathcal{N}}^\infty(p_-, p_+)$ can be approximated by $\widetilde{x} \in \widetilde{\mathcal{M}}^\lambda(p_-, p_+)$, so that in particular, when $\text{index}(p_-, \mathcal{F}) - \text{index}\,(p_+, \mathcal{F}) = 1$, there is a canonical one-to-one correspondence between $\mathcal{N}^\infty(p_-, p_+)$ and $\mathcal{M}^\lambda(p_-, p_+)$.

Since we have assumed that the restriction of (f, g) to $\mu^{-1}(0)$ is Morse-Smale, the trajectory y is transverse. Namely, the linearized operator

$$\begin{array}{rccc} \overline{D}_y: & W^{1,2}(y^* T\mu^{-1}(0)) & \to & L^2(y^* T\mu^{-1}(0)) \\ & V & \mapsto & \nabla_{y'} V + \nabla_V (\nabla f + \widetilde{\eta} \nabla \mu) \end{array} \tag{3.18}$$

is surjective and has a bounded right inverse $\overline{Q}_y$. In particular, we can choose $\overline{Q}_y$ such that

$$\text{Im}\overline{Q}_y = \left\{ V \in W^{1,2}(y^* T\mu^{-1}(0)) \mid g(V(0), W(0)) = 0 \ \forall W \in \ker \overline{D}_y \right\}. \tag{3.19}$$

For any large λ, we take our approximate solution just to be

$$\widetilde{y}(t) = (y(t), \widetilde{\eta}(y(t))). \tag{3.20}$$

Note that $y(t)$ converges to $p_\pm$ exponentially as $t \to \pm\infty$, so $\widetilde{y} \in \mathcal{B}$. Then denote the linearization of $\widetilde{S}^\lambda$ at $\widetilde{y}$ by

$$\begin{array}{rccc} \widetilde{D}_{\widetilde{y}}: & W^{1,2}(y^*TM \oplus \mathbb{R}) & \to & L^2(y^*TM \oplus \mathbb{R}) \\ & \begin{pmatrix} V \\ h \end{pmatrix} & \mapsto & \begin{pmatrix} \nabla_{y'}V + \nabla_V(\nabla f + \widetilde{\eta}\nabla\mu) + h\nabla\mu \\ h' + \lambda^2 d\mu(V) \end{pmatrix}. \end{array} \tag{3.21}$$

Now we construct a right inverse to $\widetilde{D}_{\widetilde{y}}$ out of $\overline{Q}_y$. Note that, for any $\overline{V} \in W_T(\widetilde{y})$

$$\widetilde{D}_{\widetilde{y}}(\overline{V}) = \big(\nabla_t\overline{V} + \nabla_{\overline{V}}(\nabla f + \eta\nabla\mu), \lambda^2 d\mu(\overline{V})\big) = \big(\overline{D}_y(\overline{V}), 0\big) \in \mathcal{E}_T(\widetilde{y}). \tag{3.22}$$

Hence with respect to the decomposition (3.6) and (3.7), the linearized operator can be written as

$$\widetilde{D}_{\widetilde{y}} = \begin{pmatrix} \overline{D}_y & A_y \\ 0 & D'_{\widetilde{y}} \end{pmatrix}. \tag{3.23}$$

Here A_y is given by

$$A_y(h_1\nabla\mu, h_2) = h_1 W \tag{3.24}$$

where W is a smooth tangent vector field on $\mu^{-1}(0)$. Hence there exists $c_1 > 0$ such that

$$\|A_y(h_1\nabla\mu, h_2)\|_{L_\lambda} \le c_1 \|h_1\|_{L^2} \le c_1\lambda^{-1}\|h_1\|_{W_\lambda}. \tag{3.25}$$

Now we look at the operator $D'_{\widetilde{y}}$. After trivialize the bundle $\mathbb{R}\{\nabla\mu\} \oplus \mathbb{R}$ isometrically to $\mathbb{R}^2$, we see the operator $D'_{\widetilde{y}}$ is transformed into

$$\mathfrak{D}'_{\widetilde{y}} = \frac{d}{dt} + \begin{pmatrix} a(t) & \lambda b(t) \\ \lambda b(t) & 0 \end{pmatrix} =: \frac{d}{dt} + B(t). \tag{3.26}$$

It is easy to see that, for λ large, $B(t)$ has 1 positive eigenvalue and one negative eigenvalue, both of which are bounded away from zero by $c\lambda$, where c is a constant independent of λ. Hence we have a (unique)bounded right inverse

$$Q'_{\widetilde{y}} : \mathcal{E}_L(\widetilde{y}) \to W_L(\widetilde{y}), \ \left\|Q'_{\widetilde{y}}\right\| \le c_2 \tag{3.27}$$

for some $c_2 > 0$. Then define

$$Q_{\widetilde{y}} = \begin{pmatrix} \overline{Q}_y & 0 \\ 0 & Q'_{\widetilde{y}} \end{pmatrix} \tag{3.28}$$

which serves the approximate right inverse.

Lemma 3.4. *There exists $\Lambda_0 > 0$, such that for all $\lambda > \Lambda_0$,*

$$\left\|\mathrm{Id} - \widetilde{D}_{\widetilde{y}}Q_{\widetilde{y}}\right\|_{L_\lambda} < \frac{1}{2} \tag{3.29}$$

with respect to the operator norm of the space L_λ.

Proof. Note that $\mathrm{Id} - \widetilde{D}_{\widetilde{y}}Q_{\widetilde{y}} = A_y \circ Q'_{\widetilde{y}}$ and

$$\left\|A_{\widetilde{y}}Q'_{\widetilde{y}}(h_1\nabla\mu, h_2)\right\|_{L_\lambda} \le \frac{c_1}{\lambda}\left\|Q'_{\widetilde{y}}(h_1\nabla\mu, h_2)\right\|_{W_\lambda} \le \frac{c_1c_2}{\lambda}\|(h_1\nabla\mu, h_2)\|_{L_\lambda}. \tag{3.30}$$

□

Hence $\widetilde{D}_{\widetilde{y}} Q_{\widetilde{y}}$ is invertible and a right inverse can be constructed as

$$\widetilde{Q}_{\widetilde{y}} = Q_{\widetilde{y}} \left(\widetilde{D}_{\widetilde{y}} Q_{\widetilde{y}} \right)^{-1} : L_\lambda \to W_\lambda. \tag{3.31}$$

And it is uniformly bounded by some constant which is independent of λ.

Now since we want to apply the implicit function theorem (cf. [13, Appendix A]), we need to trivialize the Banach manifold $\mathcal{B}$ and the Banach space bundle $\mathcal{E}$ locally near $\widetilde{y}$. We identify a neighborhood of $\widetilde{y}$ in $\mathcal{B}$ with a small ball (with respect to the W_λ-norm) centered at the origin of $T_{\widetilde{y}}\mathcal{B}$ as follows: using the identification $U_\epsilon \simeq \mu^{-1}(0) \times (-\epsilon, \epsilon)$, for any $\widetilde{W} = (W, a\nabla\mu, h) \in T_{\widetilde{y}}\mathcal{B}$, we identify it with the map $\rho(\widetilde{W}) = (\overline{\exp}_{\widetilde{y}} W, a, \widetilde{\eta}(\widetilde{y}) + h) \in \mu^{-1}(0) \times (-\epsilon, \epsilon) \times \mathbb{R}$ where $\overline{\exp}$ is the exponential map inside $\mu^{-1}(0)$. We trivialize $\mathcal{E}$ over such a neighborhood around $\widetilde{y}$ by parallel transport along the radial geodesics, which is denoted by $\Phi_{\widetilde{W}} : \mathcal{E}_{\rho(\widetilde{W})} \to \mathcal{E}_{\widetilde{y}}$. Then for $\widetilde{W} \in T_{\widetilde{y}}\mathcal{B}$ with $\left\| \widetilde{W} \right\|_{W_\lambda}$ small, denote by $\widetilde{D}_{\widetilde{W}} : T_{\widetilde{y}}\mathcal{B} \to \mathcal{E}_{\widetilde{y}}$ the linearization of $\Phi_{\widetilde{W}} \circ \widetilde{S}^\lambda(\rho(\widetilde{W}))$ at $\widetilde{W}$. We have

Lemma 3.5. *There exists $\epsilon_3, c_3 > 0$, independent of λ, such that for all $\widetilde{W} \in T_{\widetilde{y}}\mathcal{B}$ with $\left\| \widetilde{W} \right\|_{W_\lambda} < \epsilon_3$, one has*

$$\left\| \widetilde{D}_{\widetilde{W}} - \widetilde{D}_{\widetilde{y}} \right\| \leq c_3 \left\| \widetilde{W} \right\|_{W_\lambda}. \tag{3.32}$$

Proof. First consider the case when the $\mathbb{R}$-component of $\widetilde{W}$ is zero. We denote by $\widetilde{S}_1$ the first component of $\widetilde{S}^\lambda$, which is independent of λ. Then we see

$$\left\| D\widetilde{S}_1(\widetilde{W}) - D\widetilde{S}_1(\widetilde{y}) \right\| \leq c \left\| \widetilde{W} \right\|_{L^\infty} \tag{3.33}$$

which is standard. Now because the $\mathbb{R}$-component of $\widetilde{W}$ is zero, we have $\left\| \widetilde{W} \right\|_{L^\infty} \leq c \left\| \widetilde{W} \right\|_{W_\lambda}$ by our definition of the W_λ-norm (3.8).

The second component of $\widetilde{S}^\lambda$ is denoted by $\widetilde{S}_2^\lambda$. Then we see in this case, for any $\widetilde{V} = (V, h) \in T_{\widetilde{y}}\mathcal{B}$

$$D\widetilde{S}_2^\lambda(\widetilde{W})(\widetilde{V}) = \lambda^2 h. \tag{3.34}$$

Hence the variation of the derivative of $\widetilde{S}_2^\lambda$ is always zero.

Now we consider the case $\widetilde{W} = (0, h_0)$ with $h_0 \in W^{1,2}(\mathbb{R})$. Then for any $\widetilde{W}_0 \in T_{\widetilde{y}}\mathcal{B}$ small, by our definition of our norms (3.8) and (3.9),

$$\begin{aligned} \left\| (\widetilde{D}_{\widetilde{W}+\widetilde{W}_0} - \widetilde{D}_{\widetilde{W}_0})(V, h) \right\|_{L_\lambda} &= \| h_0 \nabla_V \nabla\mu \|_{L_\lambda} = \| h_0 \nabla_V \nabla\mu \|_{L^2} \\ &\leq c \left\| \widetilde{W} \right\|_{W_\lambda} \|V\|_{L^2} \leq c \left\| \widetilde{W} \right\|_{W_\lambda} \|(V, h)\|_{W_\lambda} \end{aligned} \tag{3.35}$$

for some $c > 0$.

Combining (3.33), (3.34) and (3.35) we obtain (3.32). □

On the other hand, we see

$$\left\|\widetilde{S}^{\lambda}(\widetilde{y})\right\|_{L_\lambda} = \lambda^{-1}\left\|\frac{d}{dt}\widetilde{\eta}(y)\right\|_{L^2} \leq c_4\lambda^{-1} \tag{3.36}$$

for some constant $c_4 > 0$. By the implicit function theorem, there exists $\epsilon_4 > 0$ such that for sufficiently large λ, there exists a unique $\widetilde{W}_\lambda(y) \in T_{\widetilde{y}}\mathcal{B}$ satisfying

$$\left\|\widetilde{W}_\lambda(y)\right\|_{W_\lambda} \leq \epsilon_4,\ \widetilde{W}_\lambda(y) \in \mathrm{Im}Q_{\widetilde{y}},\ \widetilde{S}^{\lambda}(\exp_{\widetilde{y}}\widetilde{W}_\lambda(y)) = 0. \tag{3.37}$$

Moreover, there exists $c_5 > 0$ such that

$$\left\|\widetilde{W}_\lambda(y)\right\|_{W_\lambda} \leq \frac{c_5}{\lambda}. \tag{3.38}$$

We hence define the *gluing map* (for large Λ_0) to be

$$\begin{array}{rccc} \Phi: & [\Lambda_0, +\infty) \times \widetilde{\mathcal{N}}^{\infty}(p_-, p_+) & \to & \cup_{\lambda \geq \Lambda_0}\widetilde{\mathcal{M}}^{\lambda}(p_-, p_+) \\ & (\lambda, y) & \mapsto & \exp_{\widetilde{y}}\widetilde{W}_\lambda(y). \end{array} \tag{3.39}$$

Theorem 3.6. *For each $\lambda \in [\Lambda_0, +\infty) \cap \Lambda^{reg}$, the restriction of Φ to $\{\lambda\} \times \widetilde{\mathcal{N}}^{\infty}(p_-, p_+)$ is a homeomorphism onto $\widetilde{\mathcal{M}}^{\lambda}(p_-, p_+)$.*

Proof. To complete the proof of this theorem, it remains to prove the local surjectivity of the gluing map Φ. For simplicity, we only prove for the case when the Morse indices of p_- and p_+ differ by one, which is enough for our purpose. Suppose it is not the case, then there exists a sequence $\lambda_i \to \infty$ and $\widetilde{x}_i \in \widetilde{\mathcal{M}}^{\lambda_i}(p_-, p_+)$ which converges to some $\widetilde{y} \in \widetilde{\mathcal{N}}^{\infty}(p_-, p_+)$ as described by Proposition 3.3 (with $n = 1$ and $t_{1,\nu} = 0$) but don't lie in the range of Φ. Then, consider the hyperplanes $N(y(s)) = (y'(s))^{\perp} \subset T_{y(s)}M$ for $s \in \mathbb{R}$. For each i, there exists $s_i \in \mathbb{R}$ such that $s_i \to 0$ and

$$\widetilde{x}_i(0) \in \left\{(\exp_{y(s_i)} V, \eta) \mid \eta \in \mathbb{R},\ V \in N(y(s_i)),\ \|V\| \leq \epsilon\right\}. \tag{3.40}$$

Then if we write $\widetilde{x}_i(t) = \exp_{\widetilde{y}(s_i+t)}\widetilde{W}_i(t)$, then $\widetilde{W}_i \in \mathrm{Im}Q_{\widetilde{y}(s_i+\cdot)}$ and $\left\|\widetilde{W}_i\right\|_{W_{\lambda_i}} \leq c_0\lambda_i^{-1}$ by Proposition 3.3. By the uniqueness part of the implicit function theorem, this implies that $\widetilde{x}_i = \Phi(\lambda_i, \widetilde{y}(s_i + \cdot))$, which contradicts with our assumption. □

Now Theorem 3.1 is an immediate consequence.

3.2. **Proof of Theorem 3.1 using geometric singular perturbation theory.** In the system (1.6)–(1.7), in which the time variable is t, we make the rescalings

$$\eta = \frac{\rho}{\epsilon}, \quad \lambda = \frac{1}{\epsilon}, \quad t = \epsilon\tau.$$

The system becomes

$$\frac{dx}{d\tau} = -\left(\epsilon\nabla f(x) + \rho\nabla\mu(x)\right), \tag{3.41}$$

$$\frac{d\rho}{d\tau} = -\mu(x). \tag{3.42}$$

Studying the limit $\lambda \to \infty$ in (1.6)–(1.7) is equivalent to studying the limit $\epsilon \to 0$ in (3.41)–(3.42). Setting $\epsilon = 0$ in (3.41)–(3.42), we obtain

$$\frac{dx}{d\tau} = -\rho \nabla \mu(x), \tag{3.43}$$

$$\frac{d\rho}{d\tau} = -\mu(x). \tag{3.44}$$

Let N denote the submanifold of M defined by $\mu = 0$. Then the set of equilibria of (3.43)–(3.44) is the set $\mu = \rho = 0$, i.e., $N_0 = N \times \{0\} \subset M \times \mathbb{R}$. N_0 is a compact codimension-two submanifold of $M \times \mathbb{R}$.

Choosing coordinates on M and linearizing (3.43)–(3.44) at an equilibrium $(x, 0)$, we obtain

$$\begin{pmatrix} \dot v \\ \dot\rho \end{pmatrix} = \begin{pmatrix} 0 & -\nabla\mu(x) \\ -d\mu(x) & 0 \end{pmatrix} \begin{pmatrix} v \\ \rho \end{pmatrix}. \tag{3.45}$$

Since 0 is a regular value of μ by (A3), $d\mu(x)$ and $\nabla\mu(x)$ are nonzero row and column vectors respectively. Therefore the matrix in (3.45) has rank 2, so it has at least $n-2$ zero eigenvalues. (Of course this is a consequence of the fact that we linearized (3.43)–(3.44) at a point on a manifold of equilibria of dimension $n-2$.) The other eigenvalues are $\pm\|\nabla\mu(x)\|$, where $\|v\| = \|v\|_{g(x)}$; this is shown by the following calculations, which use the fact

$$d\mu(x)v = \langle \nabla\mu(x), v\rangle_{g(x)}.$$

$$\begin{pmatrix} 0 & -\nabla\mu(x) \\ -d\mu(x) & 0 \end{pmatrix} \begin{pmatrix} \nabla\mu(x) \\ -\|\nabla\mu(x)\| \end{pmatrix} = \begin{pmatrix} \|\nabla\mu(x)\|\nabla\mu(x) \\ -\|\nabla\mu(x)\|^2 \end{pmatrix},$$

$$\begin{pmatrix} 0 & -\nabla\mu(x) \\ -d\mu(x) & 0 \end{pmatrix} \begin{pmatrix} \nabla\mu(x) \\ \|\nabla\mu(x)\| \end{pmatrix} = \begin{pmatrix} -\|\nabla\mu(x)\|\nabla\mu(x) \\ -\|\nabla\mu(x)\|^2 \end{pmatrix}.$$

Therefore $N \times \{0\}$ is a compact normally hyperbolic manifold of equilibria for (3.41)–(3.42) with $\epsilon = 0$; see Appendix A. It follows that for small $\epsilon > 0$, (3.41)–(3.42) has a normally hyperbolic invariant manifold N_ϵ near N_0.

Locally we may assume the coordinates on M are chosen so that $\mu = x_n$. Let $y = (x_1, \ldots, x_{n-1})$, so $x = (y, x_n)$. Then locally N_ϵ is parameterized by y and is given by

$$x_n = x_n(y,\epsilon) = x_n(x_1,\ldots,x_{n-1},\epsilon), \quad \rho = \rho(y,\epsilon) = \rho(x_1,\ldots,x_{n-1},\epsilon),$$

with $x_n(y,0) = \rho(y,0) = 0$. After division by ϵ, the system (3.41)–(3.42) restricted to N_ϵ is given by

$$\dot y = -\nabla_y f(y,0) + \mathcal{O}(\epsilon),$$

where $\nabla_y f(y, x_n)$ denotes the first $n-1$ components of $\nabla f(y, x_n)$.

It follows that the system (3.41)–(3.42) restricted to N_ϵ is a perturbation of the negative gradient flow of $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$. Since $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ is Morse-Smale by (A5), its negative gradient flow is structurally stable. Therefore, for small $\epsilon > 0$, the flow of (3.41)–(3.42) restricted to N_ϵ is topologically equivalent to the negative gradient flow of $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$.

An equilibrium x of the negative gradient flow of $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ corresponds, for each $\lambda > 0$, to an equilibrium (x, η) of (1.6)–(1.7), which turn corresponds to the equilibrium

$(x, \epsilon\eta)$ of (3.41)–(3.42); the latter lies in N_ϵ, which contains all complete orbits nearby. It has one higher index than that of x for the negative gradient flow of $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$. (The reason is that one of the transverse eigenvalues computed above is positive.) If the only connections between these equilibria are those in N_ϵ, the resulting Morse-Smale-Witten chain complex is the same as that of $(f|_{\mu^{-1}(0)}, g|_{\mu^{-1}(0)})$ with degree shifted by one.

To rule out the existence of other connections, note that for $\epsilon > 0$, the energy $E_\epsilon(x, \rho) = \epsilon f(x) + \rho\mu(x)$ decreases along solutions of (3.41)–(3.42). For small $\epsilon > 0$, the energy difference between two equilibria $(x_-, \epsilon\eta_-)$ and $(x_+, \epsilon\eta_+)$ of (3.41)–(3.42) is of order ϵ.

On the other hand, by the normal hyperbolicity of N_0, any sufficiently small neighborhood V of N_0 has the property that for (3.41)–(3.42) with ϵ small, a solution of (3.41)–(3.42) that starts in $V \setminus N_\epsilon$ must leave V in forward or backward time. Therefore a solution of (3.41)–(3.42) that connects two equilibria but does not lie in N_ϵ must at some time leave V through its boundary. If we can show that it must do so at a point where E_ϵ is of order one, we have a contradiction.

For a small $\alpha > 0$, we can take $V = \{(x, \rho) : |\mu(x)| < \alpha, |\rho| < \alpha, |\rho\mu(x)|) < \frac{\alpha^2}{4}\}$; see Figure 1. At points on the boundary where $|\rho\mu(x)| = \frac{\alpha^2}{4}$, $|E_\epsilon|$ is close to $\frac{\alpha^2}{4}$ for small ϵ. Thus if we can show that a solution that connects two equilibria must leave V through such a point, we are done.



FIGURE 1. The set V.

This is not true, but by making a small ϵ-dependent alteration in V, we can make it true: we replace the portions of ∂V on which $\mu = \pm\alpha$ or $\rho = \pm\alpha$ by nearby invariant surfaces, so that solutions cannot cross them. More precisely, we replace the portion of the boundary on which $\mu = \pm\alpha$ by a union of integral curves of (3.41)–(3.42) that start on the codimension-two surfaces $\mu = \pm\alpha$, $\rho = 0$, and we replace the portion of the boundary on which $\rho = \pm\alpha$ by a union of integral curves of (3.41)–(3.42) that start on the codimension-two surfaces $\mu = 0$, $\rho = \pm\alpha$. Details are left to the reader.

4. FAST-SLOW SYSTEM ASSOCIATED TO THE $\lambda \to 0$ LIMIT

Now we consider the limit of (1.6)–(1.7) as $\lambda \to 0$. In this limit, (1.6)–(1.7) is a "fast-slow system" [9]. There is one slow variable, η. In this section, we will identify the slow manifold, and study the fast flow and slow equation, under appropriate generic assumptions. (This terminology is reviewed in Appendix A.) We will also relate various Morse indices.

4.1. **The slow manifold.** The *slow manifold* for (1.6)–(1.7) is the set of equilibria for $\lambda = 0$:

$$\mathcal{C}_{\mathcal{F}} = \{(x,\eta) \in M \times \mathbb{R} \mid \nabla f(x) + \eta \nabla \mu(x) = 0\}.$$

We assume:

(A6) $\mathrm{Crit}(f) \cap \mathrm{Crit}(\mu)$ is empty.
(A7) $\nabla f + \eta \nabla \mu$ is a transverse section of $\pi^* TM$, where $\pi : M \times \mathbb{R} \to M$ is the projection.

(A7) implies that $\mathcal{C}_{\mathcal{F}}$ is a 1-dimensional smooth submanifold of $M \times \mathbb{R}$. Indeed, it is noncompact and its end at infinity is asymptotic to $(\mathrm{Crit}\mu \times [R, +\infty)) \cup (\mathrm{Crit}\mu \times (-\infty, -R])$ for R large. This can be seen as follows. If there is a sequence $(x_i, \eta_i) \in \mathcal{C}_{\mathcal{F}}$ with $\lim_{i\to\infty} |\eta_i| = \infty$, then we see that $\nabla\mu(x_i) \to 0$ and a subsequence of x_i converging to some $y \in \mathrm{Crit}\mu$. Conversely, for any $y \in \mathrm{Crit}\mu$, let $S_\epsilon(y) \subset M$ denote the sphere of radius ϵ around y. Since by (A2) y is a nondegenerate critical point of μ, we see that for small $\epsilon > 0$, the map

$$S_\epsilon(y) \to S(T_y M),\ x \mapsto \frac{\nabla\mu(x)}{\|\nabla\mu(x)\|} \tag{4.1}$$

has degree ± 1. On the other hand, by (A6) we can use local coordinates near y in which the vector field ∇f is constant. Hence there exist a unique pair of points $(x_\epsilon^\pm, \eta_\epsilon^\pm)$, with $x_\epsilon^\pm \in S_\epsilon(p)$, such that

$$(x_\epsilon^\pm, \eta_\epsilon^\pm) \in \mathcal{C}_{\mathcal{F}}. \tag{4.2}$$

We can order the two points so that $\lim_{\epsilon\to 0} \eta_\epsilon^\pm = \pm\infty$.

For fixed η, denote the function $f_\eta = f + \eta\mu : M \to \mathbb{R}$. Then consider the function

$$\begin{array}{rccc} d_{\mathcal{C}} : & \mathcal{C}_{\mathcal{F}} & \to & \mathbb{R} \\ & p = (x,\eta) & \mapsto & \det \mathrm{Hess} f_\eta(x). \end{array} \tag{4.3}$$

We assume

(A8) 0 is a regular value of $d_{\mathcal{C}}$.
(A9) For any $p = (x,\eta) \in d_{\mathcal{C}}^{-1}(0)$, $\mu(x) \neq 0$. Therefore $d_{\mathcal{C}}^{-1}(0) \cap \mathrm{Crit}(\mathcal{F}) = \emptyset$.

We denote $\mathcal{C}_{\mathcal{F}}^{sing} = d_{\mathcal{C}}^{-1}(0)$. From (A8), as $(x,\eta) \in \mathcal{C}_{\mathcal{F}}$ crosses a point in $\mathcal{C}_{\mathcal{F}}^{sing}$, one and only one of the eigenvalues of $\mathrm{Hess} f_\eta(x)$ changes its sign.

By the implicit function theorem, $\mathcal{C}_{\mathcal{F}}$ can be smoothly parameterized by η near any point of $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$ as $x = x(\eta)$. On the other hand, near $\mathcal{C}_{\mathcal{F}}^{sing}$ we have the following result.

Proposition 4.1. *Let $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}^{sing}$. We can choose local coordinates $(x_1, \ldots, x_n)$ on M such that x_p corresponds to $(0, \ldots, 0)$, and near p, $\mathcal{C}_{\mathcal{F}}$ is parameterized by x_n. Moreover, for some $c \neq 0$,*

$$\eta = \eta_p + c x_n^2 + O(x_n^3). \tag{4.4}$$

Thus $\eta|_{\mathcal{C}_{\mathcal{F}}}$ has a nondegenerate critical point at p.

This is a standard result, but we shall give a proof in Subsection 4.3.

4.2. **The fast flow.** The fast flow $\Phi_t : M \times \mathbb{R} \to M \times \mathbb{R}$ of (1.6)–(1.7) is the flow on $M \times \mathbb{R}$ determined by (1.6)–(1.7) for $\lambda = 0$. The set of equillibria of the fast flow is just $\mathcal{C}_{\mathcal{F}}$, and along the fast flow, η is constant.

If $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}$, then x_p is a critical point of the function $f_{\eta_p} : M \to \mathbb{R}$. A *fast solution* from $p_- \in \mathcal{C}_{\mathcal{F}}$ to $p_+ \in \mathcal{C}_{\mathcal{F}}$ is a solution $\widetilde{x}(t)$ of (1.6)–(1.7) for $\lambda = 0$ such that $\lim_{t\to\pm\infty} \widetilde{x}(t) = p_\pm$. A *fast orbit* from p_- to p_+ is an equivalence class of fast solutions from p_- to p_+ modulo time translation. A fast solution or orbit is *trivial* if the orbit consists of a single point.

For $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}$, x_p is a nondegenerate critical point if and only if $p \in \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$. Note that the Morse index of a nondegenerate critical point x of f_η can be defined as the number of negative eigenvalues of the matrix $\text{Hess} f_\eta(x)$. We define the Morse index of a degenerate critical point the same way. We shall use the already introduced notation $\text{index}\,(x_p, f_{\eta_p})$ to denote the Morse index in this sense of x_p as a critical point of f_{η_p}.

$\text{Index}\,(x_p, f_{\eta_p})$ is locally constant on $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$. Near $p \in \mathcal{C}_{\mathcal{F}}^{sing}$, the values of $\text{index}\,(x_p, f_{\eta_p})$ on the two branches differ by one, and $\text{index}\,(x_p, f_{\eta_p})$ is the lower of these two numbers.

4.2.1. *Transversality assumptions.* On $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$, the fast flow is normally hyperbolic. For any subset β of $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$ define

$$W^u(\beta) = \left\{ (x, \eta) \in M \times \mathbb{R} \mid \lim_{t\to-\infty} \Phi_t(x, \eta) \in \beta \right\} \tag{4.5}$$

and

$$W^s(\beta) = \left\{ (x, \eta) \in M \times \mathbb{R} \mid \lim_{t\to+\infty} \Phi_t(x, \eta) \in \beta \right\}. \tag{4.6}$$

Note that in this section we always have $\lambda = 0$, so we will not use the notation of Subsection 2.3 to specify a value of λ. If β is connected, define $\text{index}\,(\beta)$ to be $\text{index}\,(x_p, f_{\eta_p})$ for any $p = (x_p, \eta_p)$ in β.

We assume

(A10) $W^u\left(\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}\right)$ and $W^s\left(\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}\right)$ intersect transversally in $M \times \mathbb{R}$.
(A11) For each $p = (x_p, \eta_p) \in \text{Crit}(\mathcal{F})$, the pair (f_{η_p}, g) is Morse-Smale on M.

Assumption (A10) can be thought of as a weak version of the Morse-Bott-Smale transversality condition (see [1]) for a Morse-Bott function and a metric. The relation between the two assumptions is that (A11) implies that the transversality in (A10) holds near certain points, but (A10) does not imply (A11).

There are two special types of nontrivial fast orbits that play important roles in our construction. We introduce them in the remainder of this subsection.

4.2.2. *Handle-slides and cusp orbits.* By (A10) we see, if $\beta_1, \beta_2 \subset \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$ are connected, then

$$\dim\, (W^u(\beta_1) \cap W^s(\beta_2)) = \text{index}\,(\beta_1) - \text{index}\,(\beta_2) + 1. \tag{4.7}$$

In particular, when index $(\beta_1) =$ index (β_2), the dimension is one, so the intersection consists of discrete nontrivial fast orbits. We call such an orbit a *handle-slide*. If a handle-slide is contained in $M \times \{\eta\}$, we say that a handle-slide happens at η.

We assume:

(A12) The composition $\pi : \mathrm{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing} \hookrightarrow M \times \mathbb{R} \to \mathbb{R}$ is injective. Moreover, handle-slides happen at distinct values of η that are not in the image of π.

Now we look near a point $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}^{sing}$. All the eigenvalues of $\mathrm{Hess} f_{\eta_p}(x_p)$ have absolute values great than a constant $a > 0$, except for the one that changes its sign at p. Then we define $W_a^u(p)$ to be the space of maps $u : (-\infty, 0] \to M$ satisfying

1. $(u(t), \eta_p) = \Phi_t(u(0), \eta_p)$ for $t \leq 0$;
2. For sufficiently large T, for $t \in (-\infty, -T]$,

$$u(t) = \exp_{x_p} V(t), \quad V \in W_a^{1,2}((-\infty, -T], T_{x_p}M).$$

Here for any open subset $\Omega \subset \mathbb{R}$, the space $W_a^{1,2}(\Omega)$ is the space of functions f with $e^{a|t|} f \in W^{1,2}(\Omega)$.

Similarly we can define $W_a^s(p)$. By the map $u \mapsto u(0)$, $W_a^u(p)$ and $W_a^s(p)$ are naturally identified with smooth submanifolds of M, which are called the a-exponential unstable and stable manifolds of p, having dimensions

$$\dim W_a^u(p) = \mathrm{index}\,(x_p, f_{\eta_p}),\ \dim W_a^s(p) = n - \mathrm{index}\,(x_p, f_{\eta_p}) - 1. \tag{4.8}$$

Let $\mathring{W}^u(p) = W^u(p) \setminus W_a^u(p)$ and $\mathring{W}^s(p) = W^s(p) \setminus W_a^s(p)$. These manifolds are unions of orbits that approach p polynomially rather than exponentially. They are submanifolds of $M \times \{\eta_p\}$ of dimension $\dim W_a^u(p) + 1$ and $\dim W_a^s(p) + 1$. We assume

(A13) For each $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}^{sing}$ and $q = (x_q, \eta_q) \in \mathcal{C}_{\mathcal{F}}$ with $\eta_p = \eta_q = \eta$, $W_a^u(x_p)$ and $W^s(x_q)$ (respectively $W_a^s(x_p)$ and $W^u(x_q)$) intersect transversely in $M \times \{\eta\}$, $\mathring{W}^u(x_p)$ and $W^s(x_q)$ (respectively $\mathring{W}^s(x_p)$ and $W^u(x_q)$) intersect transversely in $M \times \{\eta\}$.

The reader may refer to [5] and [4] for this transversality in the case of Lagrangian intersections.

(A13) implies that for $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}^{sing}$ and $q = (x_q, \eta_q) \in \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$, $W_a^u(p) \cap W^s(q) = \emptyset$ if $\mathrm{index}\,(x_q, f_{\eta_q}) \geq \mathrm{index}\,(x_p, f_{\eta_p}) - 1$; and $W^u(q) \cap W_a^s(p) = \emptyset$ if $\mathrm{index}\,(x_p, f_{\eta_p}) \geq \mathrm{index}\,(x_q, f_{\eta_q}) - 2$. For example, $W^u(q) \cap W_a^s(p) = \emptyset$ if

$$\begin{aligned} 0 \geq \dim W^u(q) + \dim W_a^s(p) - (n+1) &= \mathrm{index}\,(x_q, f_{\eta_q}) + (n - \mathrm{index}\,(x_p, f_{\eta_p}) - 1) - (n+1) \\ &= \mathrm{index}\,(x_q, f_{\eta_q}) - \mathrm{index}\,(x_p, f_{\eta_p}) - 2. \end{aligned} \tag{4.9}$$

When the inequalities are equalities, $W^u(p) \cap W^s(q) = \mathring{W}^u(p) \cap W^s(q)$ and $W^u(q) \cap W^s(p) = W^u(q) \cap \mathring{W}^s(p))$ consist of isolated orbits corresponding to solutions that approach p polynomially. We call such orbits *cusp orbits*; parametrized ones are *cusp solutions*. An argument similar to one in [4] shows that between p and q there are only finitely many cusp orbits.

4.2.3. *Short fast orbits.* By Proposition 4.1, near $p \in \mathcal{C}_{\mathcal{F}}^{sing}$, one can parametrize $\mathcal{C}_{\mathcal{F}}$ by $\gamma_p : (-\epsilon, \epsilon) \to \mathcal{C}_{\mathcal{F}}$ such that

$$\gamma_p(0) = p, \ \eta(t) = \eta_p \pm s^2. \tag{4.10}$$

We choose the orientation of γ_p such that $\operatorname{index}(\gamma_p((-\epsilon, 0))) = \operatorname{index}(\gamma_p((0, \epsilon))) + 1$. For ϵ small enough there exists a unique such parametrization γ and we call γ the *canonical parametrization* near p. Then, for ϵ small enough, there is a unique trajectory of the flow of $-\nabla f_{\eta(s)}$ from $\gamma(-s)$ to $\gamma(s)$ and they are all contained in a small neighborhood of x_p.

Lemma 4.2. *There exists $\epsilon_0 > 0$ such that for all $p \in \mathcal{C}_{\mathcal{F}}^{sing}$ and for $s \in (0, \epsilon_0)$ there exists a unique trajectory of the fast flow from $\gamma_p(-s)$ to $\gamma_p(s)$, where $\gamma_p : (-\epsilon_0, \epsilon_0) \to \mathcal{C}_{\mathcal{F}}$ is the canonical parametrization of $\mathcal{C}_{\mathcal{F}}$ near p.*

If $0 < s \leq \epsilon < \epsilon_0$, then we call a fast orbit from $\gamma_p(-s)$ to $\gamma_p(s)$ an ϵ-*short orbit.*

Lemma 4.3. *For any ϵ small enough, there exists $\delta = \delta(\epsilon) > 0$ such that, for all $\eta \in \mathbb{R}$, any trajectories of the flow of $-\nabla f_\eta$ which is not an ϵ-short trajectory has energy no less than δ.*

Proof. Denote $\mathcal{C}_{\mathcal{F}}^{sing,\epsilon} = \cup_{p \in \mathcal{C}_{\mathcal{F}}^{sing}} \gamma_p((-\epsilon, \epsilon))$. Because we have assumed that the map $\mathcal{C}_{\mathcal{F}}^{sing} \to M \times \mathbb{R} \to \mathbb{R}$ is injective, any fast trajectory both of whose beginning and end lie in $\mathcal{C}_{\mathcal{F}}^{sing,\epsilon}$ much be an ϵ-short trajectory. So suppose the lemma doesn't hold, then there exists $\epsilon > 0$ and a sequence of nontrivial fast trajectories $(y_k, \eta_k) \subset M \times \{\eta_k\}$ such that

$$\lim_{k \to \infty} f_{\eta_k}(y_k(-\infty)) - f_{\eta_k}(y_k(+\infty)) = 0 \tag{4.11}$$

and they are not ϵ-short ones. Hence without loss of generality, we may assume that for all k, $y_k(-\infty) \notin \mathcal{C}_{\mathcal{F}}^{sing,\epsilon}$.

(1) $\lim_{k \to \infty} \eta_k = \pm\infty$, which implies that $y_k(\pm\infty)$ converges to points in $\operatorname{Crit}(\mu)$. However, if we rescale the orbit by $z_k(t) = y_k(\eta_k^{-1} t)$, then z_k converges to a (possibly broken) nontrivial trajectory of the flow of $-\nabla \mu$, which contradicts with (4.11).

(2) $\lim_{k \to \infty} \eta_k = \eta_\infty \in \mathbb{R}$. Then a subsequence of y_k converges to a broken trajectory of f_{η_∞}, which must be a constant one. Hence

$$\lim_{k \to \infty} (y_k(\pm\infty)) =: y_\infty \in \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing,\epsilon}. \tag{4.12}$$

So for k large enough, $y_k(\pm\infty)$ lie in the same connected component of $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing,\frac{\epsilon}{2}}$. This is impossible because on each such component, distinct points have distinct values of η.

□

4.3. **The slow equation.** On $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$, which is parameterized by η, the slow equation is given by restricting (1.7) to $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$ and dividing by λ^2:

$$\eta' = -\mu(x(\eta)). \tag{4.13}$$

Thus η' changes sign only when $\mu = 0$. We have

$$\operatorname{Crit}(\mathcal{F}) = \{(x, \eta) \in \mathcal{C}_{\mathcal{F}} \mid \mu(x) = 0\}.$$

Thus η' changes sign only at points of $\mathrm{Crit}(\mathcal{F})$. By (A9), η' does not change sign at points of $\mathcal{C}_{\mathcal{F}}^{sing}$.

Proposition 4.4. *Equilibria of (4.13) are hyperbolic.*

Proof. Let $p = (x_p, \eta_p)$ be an equilibrium of (4.13). Then $\mu(x_p) = 0$, so by (A3), $d\mu(x_p) \neq 0$. Hence we can choose local coordinates $(x_1, \ldots, x_n)$ near x_p such that $\mu(x_1, \ldots, x_n) = x_n$. Near p, $\mathcal{C}_{\mathcal{F}}$ can be defined without reference to the metric g by the equations

$$\frac{\partial f}{\partial x_i} + \eta\delta_{nj}, \; j = 1, \ldots, n; \quad \delta_{ij} = \text{Kronecker delta}. \tag{4.14}$$

Let $e_n = \begin{pmatrix} 0 & \ldots & 0 & 1 \end{pmatrix}^T$, the nth standard unit basis vector in $\mathbb{R}^n$. Then the derivative of the system (4.14) is the $n \times (n+1)$ matrix

$$\begin{pmatrix} \mathrm{Hess} f(x) & e_n \end{pmatrix} = \begin{pmatrix} A & b & 0 \\ b^T & c & 1 \end{pmatrix}, \tag{4.15}$$

where A is an $(n-1) \times (n-1)$ matrix, and $(b^T, c) = \left(\frac{\partial^2 f}{\partial x_1 \partial x_n}, \cdots, \frac{\partial^2 f}{\partial x_{n-1} \partial x_n}, \frac{\partial^2 f}{\partial x_n^2} \right)$.

Since x_p is a critical point of $f|_{\mu^{-1}(0)}$, by (A5), A is invertible. The matrix (4.15) has full rank; a nonzero vector in its kernel is tangent at p to $\mathcal{C}_{\mathcal{F}}$. Since $p \in \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$, such a vector has nonzero η-component, and one can check that it has nonzero x_n-component. Therefore near p, when $\mathcal{C}_{\mathcal{F}}$ is parameterized by η, we have $x(\eta) = (y(\eta), x_n(\eta))$ with

$$x_n(\eta) = a(\eta - \eta_p) + \mathcal{O}(\eta - \eta_p)^2, \quad a \neq 0.$$

But then (4.13) becomes

$$\eta' = a(\eta - \eta_p) + \mathcal{O}(\eta - \eta_p)^2,$$

which proves the result. □

Next we discuss the slow equation near points of $\mathcal{C}_{\mathcal{F}}^{sing}$. Recall that if p_0 is an equilibrium of $\dot{p} = h(p)$, h is C^s, $Dh(p_0)$ has m eigenvalues (counting multiplicity) with real part 0, and E is the corresponding m-dimensional invariant subspace of $\dot{v} = Dh(p_0)v$, then there is a C^s invariant manifold through p_0 and tangent there to E, called the *center manifold* of $\dot{p} = h(p)$ at p_0.

Proposition 4.5. *Let $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}^{sing}$. We can choose local coordinates $(x_1, \ldots, x_n)$ on M near x_p, with x_p corresponding to 0, in which*

$$D\nabla f(0) = \begin{pmatrix} A & 0 \\ 0 & 0 \end{pmatrix}, \quad A \text{ an invertible symmetric } (n-1) \times (n-1) \text{ matrix.} \tag{4.16}$$

The center manifold of (1.6)–(1.7) at p is parameterized by (x_n, η). There are numbers $c \neq 0$ and $d \neq 0$ such that the system (1.6)–(1.7), restricted to the center manifold, is

$$x_n' = c(\eta - \eta_p) + dx_n^2 + \ldots, \tag{4.17}$$

$$\eta' = -\lambda^2(\mu(x_p) + \ldots), \tag{4.18}$$

where … indicates terms of higher order (and, in (4.17), other second-order terms). Therefore near p, $\mathcal{C}_{\mathcal{F}}$ is parameterize by x_n, and $\eta|\mathcal{C}_{\mathcal{F}}$ has a nondegenerate critical point at p.

Proof. (A7) implies that we can choose local coordinates $(x_1, \dots, x_n)$ on M near x_p, with x_p corresponding to 0, in which (4.16) holds. In these coordinates, $\mu = \mu(x_p) + a^T x + \dots$ with $a \in \mathbb{R}^n$. Write $x = (y, x_n)$ with $y \in \mathbb{R}^{n-1}$, and write $a = (b, c)$ with $b \in \mathbb{R}^{n-1}$ and $c \in \mathbb{R}$. Near p the system (1.6)–(1.7) becomes

$$y' = Ay + b(\eta - \eta_p) + \dots, \tag{4.19}$$

$$x_n' = c(\eta - \eta_p) + dx_n^2 + \dots, \tag{4.20}$$

$$\eta' = -\lambda^2(\mu(x_p) + \dots), \tag{4.21}$$

where …indicates higher-order terms (and, in (4.20), other second-order terms). (A7) implies that $c \neq 0$, and (A8) implies that $d \neq 0$. The linearization of (4.19)–(4.21) at $(y, x_n, \eta) = (0, 0, \eta_p)$ is

$$\begin{pmatrix} y' \\ x_n' \\ \eta' \end{pmatrix} \begin{pmatrix} A & 0 & b \\ 0 & 0 & c \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} y \\ x_n \\ \eta - \eta_p \end{pmatrix}.$$

The matrix has a two-dimensional generalized eigenspace for the eigenvalue 0 that is tangent to the center manifold of (4.19)–(4.19) at p. The center manifold is parameterized by (x_n, η), and to lowest order it is given by

$$y = -A^{-1}b(\eta - \eta_p)$$

The restriction of (4.19)–(4.21) to the center manifold is therefore given by (4.17)–(4.18). ☐

Proposition 4.1 is of course just a partial restatement of Proposition 4.5.

4.4. **Three indices.** In this subsection we describe the relation between three different Morse indices:

1. For any $p = (x_p, \eta_p) \in \mathcal{C}_{\mathcal{F}}$, we have $\text{index}(x_p, f_{\eta_p})$, used in Subsection 4.2.
2. For $p = (x_p, \eta_p) \in \text{Crit}(\mathcal{F})$, we can consider $\text{index}\,(p, \mathcal{F})$.
3. If $p = (x_p, \eta_p) \in \text{Crit}(\mathcal{F})$, then $x_p \in \text{Crit}(f|_{\mu^{-1}(0)})$, and we can consider $\text{index}\,(x_p, f|_{\mu^{-1}(0)})$.

We have already seen in Lemma 2.1 that $\text{index}(p, \mathcal{F}) = \text{index}(x_p, f|_{\mu^{-1}(0)}) + 1$ for $p = (x_p, \eta_p) \in \text{Crit}(\mathcal{F})$. The best way to see the relation of the different indices is to look at the corresponding unstable manifolds.

Proposition 4.6. *Let* $p = (x_p, \eta_p) \in \text{Crit}(\mathcal{F})$.

1. *If p is a repeller of the slow equation, then* $\text{index}\,(p, \mathcal{F}) = \text{index}(x_p, f_{\eta_p}) + 1$.
2. *If p is an attractor of the slow equation, then* $\text{index}\,(p, \mathcal{F}) = \text{index}(x_p, f_{\eta_p})$.

Proof. If p is a repeller of the slow equation, then for small $\lambda > 0$, the equilibrium p of (1.6)–1.7) has one more positive eigenvalue than the equilibrium x_p of f_{η_p}. Since the index of a critical point of a Morse function h is the number of positive eigenvalues of the corresponding equilibrium of ∇h for any metric, (1) follows. (2) is similar. ☐

5. ADIABATIC LIMIT $\lambda \to 0$

5.1. **Slow solutions, fast solutions, and fast-slow solutions.**

Let $p_-, p_+ \in \mathcal{C}_{\mathcal{F}}$.

Recall that a nontrivial fast solution from p_- to p_+ is a nonconstant solution $\widetilde{x}(t)$, $-\infty < t < \infty$, of (1.6)–(1.7) for $\lambda = 0$, such that $\lim_{t\to\pm\infty} \widetilde{x}(t) = p_\pm$.

Let $(x(\eta), \eta)$ parameterize the closure of a component of $\mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$. Let $I(\alpha_-, \alpha_+) \subset \mathbb{R}$ be the closed interval from α_- to α_+, with $\alpha_- \leq \alpha_+$; we allow $\alpha_- = -\infty$, $\alpha_+ = +\infty$, and $\alpha_- = \alpha_+ \in \mathbb{R}$. Let $\eta(t)$, t in the interior of $I(\alpha_-, \alpha_+)$, be a solution of $\eta' = -\mu(x(\eta))$. If α_- (respectively α_+) is finite, we extend $\eta(t)$ continuously to α_- (respectively α_+). Let $p(t) = (x(\eta(t)), \eta(t))$, $t \in I(\alpha_-, \alpha_+)$, and let $p_\pm \in \mathcal{C}_{\mathcal{F}}$. Then $p(t)$ is a *slow solution from* p_- *to* p_+ (for short, a *slow solution*) provided:

1. if α_- (respectively α_+) is finite, then $p(\alpha_-) = p_-$ (respectively $p(\alpha_+) = p_+$);
2. if $\alpha_- = -\infty$ (respectively $\alpha_+ = +\infty$), then $\lim_{t\to-\infty} p(t) = p_-$ (respectively $\lim_{t\to+\infty} p(t) = p_+$).

A slow solution or its orbit is *trivial* if the orbit is a single point in $\mathcal{C}_{\mathcal{F}}$.

A *fast-slow solution from* p_- *to* p_+ of (1.6)–(1.7) (for short, a *fast-slow solution*) is a sequence

$$\mathscr{X} = (p_0, \sigma_1, p_1, \sigma_2, p_2, \ldots, p_{n-1}, \sigma_n, p_n) \tag{5.1}$$

such that:

1. $p_0 = p_-$, $p_n = p_+$.
2. Each $p_i \in \mathcal{C}_{\mathcal{F}}$.
3. Each σ_i is a nontrivial fast solution or a slow solution from p_{i-1} to p_i. Trivial slow solutions are allowed, but σ_1 and σ_n are not allowed to be trivial slow solutions.
4. Either (i) σ_i is slow for i even and fast for i odd, or (ii) σ_i is slow for i odd and fast for i even.

Trivial slow solutions are allowed to deal with the possibility that a fast solution to p is followed by a fast solution from p.

An orbit (fast orbit, slow orbit, or fast-slow orbit) is an equivalence class of solutions obtained by forgetting the parametrization (but remembering the orientation in which $\mathcal{F}$ decreases). In particular, a fast-slow orbit is denoted by $[\mathscr{X}]$ for $\mathscr{X}$ as in (5.1).

Remark 5.1. A fast-slow orbit defined above is similar to a “flow line with cascades” in Morse-Bott theory (see [6]), where the fast solutions correspond to “cascades”. The difference is that the slow manifold $\mathcal{C}_{\mathcal{F}}$ is not everywhere normally hyperbolic, and we can have $p_i \in \mathcal{C}_{\mathcal{F}}^{sing}$.

In the remainder of this subsection, we will discuss some properties of fast-slow orbits from $p_- \in \mathcal{C}_{\mathcal{F}}$ to $p_+ \in \mathcal{C}_{\mathcal{F}}$. We are particularly interested in the case $p_\pm \in \mathrm{Crit}\,(\mathcal{F})$ and $\mathrm{index}(p_-, \mathcal{F}) - \mathrm{index}(p_+, \mathcal{F}) = 1$.

The following proposition is obvious.

Proposition 5.2. *Consider a nontrivial slow solution* $p(t)$, $t \in I(\alpha_-, \alpha_+)$, *from* p_- *to* p_+.

1. *If* $p_- \in \mathcal{C}_{\mathcal{F}}^{sing}$, *then either (i) for all* $t \in (\alpha_-, \alpha_+)$, $\mathrm{index}\,(x_{p(t)}, f_{\eta_{p(t)}}) = \mathrm{index}\,(x_-, f_{\eta_-})$, *or (ii) for all* $t \in (\alpha_-, \alpha_+)$, $\mathrm{index}\,(x_{p(t)}, f_{\eta_{p(t)}}) = \mathrm{index}\,(x_-, f_{\eta_-}) + 1$.

(2) *If* $p_+ \in \mathcal{C}_{\mathcal{F}}^{sing}$*, then either (i) for all* $t \in (\alpha_-, \alpha_+)$, $\operatorname{index}(x_{p(t)}, f_{\eta_{p(t)}}) = \operatorname{index}(x_+, f_{\eta_+})$, *or (ii) for all* $t \in (\alpha_-, \alpha_+)$, $\operatorname{index}(x_{p(t)}, f_{\eta_{p(t)}}) = \operatorname{index}(x_+, f_{\eta_+}) + 1$.

A slow solution $p(t)$ for $t \in I(\alpha_-, \alpha_+)$ is *regular* if it is nontrivial and satisfies:

1. If $p_- \in \mathcal{C}_{\mathcal{F}}^{sing}$, then for all $t \in (\alpha_-, \alpha_+)$, $\operatorname{index}(x_{p(t)}, f_{\eta_{p(t)}}) = \operatorname{index}(x_-, f_{\eta_-}) + 1$.
2. If $p_+ \in \mathcal{C}_{\mathcal{F}}^{sing}$, then for all $t \in (\alpha_-, \alpha_+)$, $\operatorname{index}(x_{p(t)}, f_{\eta_{p(t)}}) = \operatorname{index}(x_+, f_{\eta_+})$.

A slow orbit is regular if it is the orbit of a regular slow solution.

See Figure 2.



FIGURE 2. (a) Direction of slow flow is downward. The thick curve is a regular slow orbit that starts at p_-. Points on it have index one greater than p_-. (b) Direction of slow flow is downward. The thick curve is a regular slow orbit that ends at p_+. Points on it have the same index as p_+.

Proposition 5.3. *Let* γ *be either a slow or fast solution from* $p_- = (x_-, \eta_-)$ *to* $p_+ = (x_+, \eta_+)$.

(1) *If* γ *is a regular slow solution, then:*
    (a) *If* $p_- \in \mathcal{C}_{\mathcal{F}}^{sing}$*, then* $\operatorname{index}(x_+, f_{\eta_+}) = \operatorname{index}(x_-, f_{\eta_-}) + 1$.
    (b) *Otherwise* $\operatorname{index}(x_+, f_{\eta_+}) = \operatorname{index}(x_-, f_{\eta_-})$

(2) *If* γ *is a nontrivial fast solution (so* $\eta_+ = \eta_-$*), then:*
    (a) *At most one of* $p_\pm$ *is in* $\operatorname{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$.
    (b) $\operatorname{index}(x_+, f_{\eta_+}) \leq \operatorname{index}(x_-, f_{\eta_-})$.
    (c) *If* $p_+ \in \operatorname{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$ *or* $p_- \in \operatorname{Crit}(\mathcal{F})$*, then* $\operatorname{index}(x_+, f_{\eta_+}) \leq \operatorname{index}(x_-, f_{\eta_-}) - 1$.

Proof. The result for slow solutions is immediate from their definition.

For fast solutions, (2)(a) follows from assumption (A12)

Let $p(t)$ be a fast solution, let $\operatorname{index}(x_-, f_{\eta_-}) = k$, and $\operatorname{index}(x_+, f_{\eta_+}) = l$. (Of course $\eta_+ = \eta_-$.)

If $p_+ \in \operatorname{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$, then (A11) or (A13) implies that $W^u(p_-)$ and $W^s(p_+)$ intersect transversally within $M \times \{\eta_-\}$. We have $\dim W^u(p_-) = k$, and $\dim W^s(p_+) = n - l$, so the intersection is nonempty provided $k + (n - l) \geq n + 1$, i.e., $l \leq k - 1$.

If $p_- \in \operatorname{Crit}(\mathcal{F})$, the same argument applies. However, if $p_- \in \mathcal{C}_{\mathcal{F}}^{sing}$, then $\dim W^u(p_-) = k + 1$. We again have $\dim W^s(p_+) = n - l$, so the intersection is nonempty provided $(k + 1) + (n - l) \geq n + 1$, i.e., $l \leq k$.

If neither p_- nor p_+ is in $\mathrm{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$, then we do not know that $W^u(p_-)$ and $W^s(p_+)$ intersect transversally within $M \times \{\eta_-\}$. However, by (A10), $(k+1) + (n-l+1) \geq n+2$, so $l \leq k$. $\square$

Proposition 5.4. *Consider a fast-slow solution* $\mathscr{X} = (p_0, \sigma_1, p_1, \sigma_2, p_2, \dots, p_{n-1}, \sigma_n, p_n)$ *from* $p_- = (x_-, \eta_-)$ *to* $p_+ = (x_+, \eta_+)$, *with neither in* $\mathcal{C}_{\mathcal{F}}^{sing}$ *and* $\mathrm{index}\,(x_+, f_{\eta_+}) \geq \mathrm{index}(x_-, f_{\eta_-}) = k$. *Then*

1. *For all* $i \geq 1$, $\mathrm{index}\,(x_{p_i}, f_{p_i}) = k$, *except for the case where* $p_i \in \mathcal{C}_{\mathcal{F}}^{sing}$ *and* σ_i *is a fast solution, in which case* $\mathrm{index}\,(x_{p_i}, f_{p_i}) = k-1$. *In particular,* $\mathrm{index}\,(x_+, f_{\eta_+}) = k$.
2. *All slow solutions appearing in* $\mathscr{X}$ *are regular. In particular, they are nontrivial.*
3. *If* $p_i \in \mathrm{Crit}(\mathcal{F})$, *then* $i = 0$ *or* $i = n$. *In the first case,* $p_0 = p_- \in \mathrm{Crit}^-(\mathcal{F})$ *and* σ_1 *is slow; in the second case,* $p_n = p_+ \in \mathrm{Crit}^+(\mathcal{F})$ *and* σ_n *is slow.*
4. *The fast solutions are handle-slides or cusp solutions.*

Proof. Along a fast-slow solution, $\mathrm{index}\,(x_{p_i}, f_{p_i}) \leq \mathrm{index}\,(x_{p_{i-1}}, f_{p_{i-1}})$, except along a regular slow solution for which $p_{i-1} \in \mathcal{C}_{\mathcal{F}}^{sing}$ and $p_i \in \mathcal{C}_{\mathcal{F}} \setminus \mathcal{C}_{\mathcal{F}}^{sing}$; in this case $\mathrm{index}\,(x_{p_i}, f_{p_i}) = \mathrm{index}\,(x_{p_{i-1}}, f_{p_{i-1}}) + 1$. However, the index drops along the fast solution from p_{i-2} to p_{i-1}. Thus (1) holds; also, nontrivial slow solutions must be regular and (3) must hold, otherwise we would have $\mathrm{index}\,(x_+, f_{\eta_+}) < \mathrm{index}(x_-, f_{\eta_-})$. (3) rules out fast solutions that start or end in $\mathrm{Crit}(\mathcal{F})$; any remaining fast solutions are handle-slides or cusp solutions, so (4) holds. Since handle-slides and cusp solutions occur at different values of η, slow solutions must be nontrivial, which completes the proof of (2). $\square$

Now we describe fast-slow solutions between two critical points whose indices differ by 1. Recall from Subsection 2.1 that $\mathrm{Crit}_k(\mathcal{F})$ denotes the set of index k critical points of $\mathcal{F}$. Let $\mathrm{Crit}_k^+(\mathcal{F})$ (respectively $\mathrm{Crit}_k^-(\mathcal{F})$) denote the set of equilibria in $\mathrm{Crit}_k(\mathcal{F})$ that are stable (respectively unstable) equilibria of the slow equation. Let $\mathrm{Crit}^{\pm}(\mathcal{F}) = \cup_{k \geq 0} \mathrm{Crit}_k^{\pm}(\mathcal{F})$.

Proposition 5.5. *Let* $p_\pm = (x_\pm, \eta_\pm) \in \mathrm{Crit}(\mathcal{F})$ *with* $p_- \in \mathrm{Crit}_k(\mathcal{F})$, $p_+ \in \mathrm{Crit}_{k-1}(\mathcal{F})$. *A fast-slow solution* $\mathscr{X} = (p_0, \sigma_1, p_1, \sigma_2, p_2, \dots, p_{n-1}, \sigma_n, p_n)$ *from* p_- *to* p_+ *has the following properties.*

1. *If* σ_i *is a fast solution from* $p_{i-1} = (x_{p_{i-1}}, \eta_{p_{i-1}})$ *to* $p_i = (x_{p_i}, \eta_{p_i})$ *(of course* $\eta_{p_i} = \eta_{p_{i-1}}$*), then* $\mathrm{index}\,(x_{p_i}, f_{\eta_{p_i}}) = \mathrm{index}\,(x_{p_{i-1}}, f_{\eta_{p_{i-1}}})$, *unless:*
    - (a) $p_i \in \mathcal{C}_{\mathcal{F}}^{sing}$, *in which case* $\mathrm{index}\,(x_{p_i}, f_{\eta_{p_i}}) = \mathrm{index}\,(x_{p_{i-1}}, f_{\eta_{p_{i-1}}}) - 1$.
    - (b) $i = 1$, $p_0 = p_- \in \mathrm{Crit}_k^+(\mathcal{F})$, *in which case* σ_1 *is fast (so* $\eta_{p_1} = \eta_p$*) and* $\mathrm{index}\,(x_{p_1}, f_{\eta_{p_1}}) = \mathrm{index}\,(x_-, f_{\eta_-}) - 1$.
    - (c) $i = n$, $p_n = p_+ \in \mathrm{Crit}_{k-1}^-(\mathcal{F})$, *in which case* σ_n *is fast (so* $\eta_q = \eta_{p_{n-1}}$*) and* $\mathrm{index}\,(x_+, f_{\eta_+}) = \mathrm{index}\,(x_{p_{n-1}}, f_{\eta_{p_{n-1}}}) - 1$.
2. *All slow solutions appearing in* $\mathscr{X}$ *are regular. In particular, they are nontrivial.*
3. *For* $i = 1, \dots, n-1$, $p_i \notin \mathrm{Crit}(\mathcal{F})$.
4. *The fast solutions in* $\mathscr{X}$ *are handle-slides or cusp solutions, except for* σ_1 *when* $p_- \in \mathrm{Crit}_k^+(\mathcal{F})$ *and* σ_n *when* $p_+ \in \mathrm{Crit}_{k-1}^-(\mathcal{F})$.

In addition:

(I) *Suppose* $p_- \in \mathrm{Crit}_k^+(\mathcal{F})$ *and* $p_+ \in \mathrm{Crit}_{k-1}^+(\mathcal{F})$*. Then the odd* σ_i *are fast, and* n *is even.*
(II) *Suppose* $p_- \in \mathrm{Crit}_k^+(\mathcal{F})$ *and* $p_+ \in \mathrm{Crit}_{k-1}^-(\mathcal{F})$*. Then the odd* σ_i *are fast, and* n *is odd.*
(III) *Suppose* $p_- \in \mathrm{Crit}_k^-(\mathcal{F})$ *and* $p_+ \in \mathrm{Crit}_{k-1}^+(\mathcal{F})$*. Then the odd* σ_i *are slow, and* n *is odd.*
(IV) *Suppose* $p_- \in \mathrm{Crit}_k^-(\mathcal{F})$ *and* $p_+ \in \mathrm{Crit}_{k-1}^-(\mathcal{F})$*. Then the odd* σ_i *are slow, and* n *is even.*

Proof. We prove (1)–(4) separately for the cases (I) and (II). The proofs for cases (III) and (IV) are similar so we omit them. We remark that in one of the cases, case (III), we can have $n = 1$, which means $\mathscr{X}$ may contain a single slow solution and no fast solutions.

Proof in case (I). In this case, $\mathrm{index}\,(x_-, f_{\eta_-}) = k$ and $\mathrm{index}\,(x_+, f_{\eta_+}) = k-1$. Since p_- is an attractor for the slow equation, the first solution σ_1 must be fast, so the odd σ_i are fast. By Proposition 5.3 (2), $p_1 \notin \mathrm{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$ and $\mathrm{index}\,(x_{p_1}, f_{\eta_{p_1}}) \le k-1$. Consider the portion of the fast-slow solution from p_1 to q. Then Proposition 5.4 implies that $\mathrm{index}\,(x_{p_1}, f_{\eta_{p_1}}) = k-1$, conclusion (2) of Proposition 5.5 holds, and $p_i \notin \mathrm{Crit}^{\pm}(\mathcal{F})$ for $i = 2, \dots, n-1$ so (3) holds. Proposition 5.4 also implies that all fast solutions except the first are handle-slides or cusp solutions, so (4) holds.

Since p_+ is an attractor for the slow equation, *a priori* the last solution could be slow or fast. However, if the last solution were fast, from Proposition 5.3 (2) we would have $\mathrm{index}\,(x_+, f_{\eta_+}) < \mathrm{index}\,(x_{p_{n-1}}, f_{\eta_{p_{n-1}}}) = k-1$, contradiction. Therefore the last solution is slow, so n is even. Then (1) and (4) follow from Proposition 5.4.

Proof in case (II). As with case (I), $\mathrm{index}\,(x_-, f_{\eta_-}) = k$; σ_1 must be fast, so the odd σ_i are fast; $p_1 \notin \mathrm{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$; and $\mathrm{index}\,(x_{p_1}, f_{\eta_{p_1}}) \le k-1$. Since p_+ is a repeller for the slow equation, $\mathrm{index}\,(x_+, f_{\eta_+}) = k-2$, and the last solution must be fast. Then $p_{n-1} \notin \mathrm{Crit}(\mathcal{F}) \cup \mathcal{C}_{\mathcal{F}}^{sing}$, and by Proposition 5.3 (2), $\mathrm{index}\,(x_{p_{n-1}}, f_{\eta_{p_{n-1}}}) \ge k-1$. As in the previous argument, for the portion of the fast-slow solution from p_1 to p_{n-1}, apply Proposition 5.4; we obtain (1)–(4). Since the first and last solutions are fast, n is odd. ☐

5.2. The chain complex.

Now let $p, q \in \mathrm{Crit}(\mathcal{F})$. We denote by $\widetilde{\mathcal{N}}^0(p,q)$ the space of all fast-slow solutions from p to q and $\mathcal{N}^0(p,q)$ the space of all fast-slow orbits from p to q.

Proposition 5.6. *If* $p, q \in \mathrm{Crit}(\mathcal{F})$ *and* $\mathrm{index}\,(q, \mathcal{F}) = \mathrm{index}\,(p, \mathcal{F}) - 1$*, then* $\#\mathcal{N}^0(p,q) < +\infty$*.*

Proof. By Proposition 5.5, we only need to show that there are finitely many handle-slides or cusp orbits contained in $\mathcal{F}^{-1}\left([\mathcal{F}(q), \mathcal{F}(p)]\right)$. But this follows from our description of $\mathcal{C}_{\mathcal{F}}$ at the beginning of Section 4 and the transversality assumption (A10), (A11), (A13). ☐

Now we define a chain complex $(\mathcal{C}^0, \partial^0)$ of $\mathbb{Z}_2$-modules. It has the same generators and gradings as $\mathcal{C}^\lambda$. Its boundary operator $\partial^0 : \mathcal{C}_k^0 \to \mathcal{C}_{k-1}^0$ is defined by

$$\langle \partial^0 p, q \rangle = \#\mathcal{N}^0(p,q) \bmod 2. \tag{5.2}$$

The main theorem of this section and the essentially new thing we constructed in this paper is:

Theorem 5.7. *$(\mathcal{C}^0, \partial^0)$ is a chain complex, i.e., $\partial^0 \circ \partial^0 = 0$. Moreover, the homology $H_k(\mathcal{C}^0, \partial^0)$ is canonically isomorphic to $H_{k-1}(\mu^{-1}(0), \mathbb{Z}_2)$.*

The proof of this theorem is carried out in the following way. Let $p, q \in \mathrm{Crit}(\mathcal{F})$. First we prove that for small $\lambda > 0$, any orbit in $\mathcal{M}^\lambda(p,q)$ is close to a fast-slow orbit in $\mathcal{N}^0(p,q)$ (the compactness theorem, Theorem 5.9). Then we restrict to the case $\mathrm{index}(p,\mathcal{F}) - \mathrm{index}(q,\mathcal{F}) = 1$ and show that for each small $\lambda > 0$, each orbit in $\mathcal{N}^0(p,q)$ is close to exactly one orbit in $\mathcal{M}^\lambda(p,q)$ (the gluing theorem, Theorem 5.14). The resulting one-to-one correspondence implies that $(\mathcal{C}^0, \partial^0)$ is isomorphic to $(\mathcal{C}^\lambda, \partial^\lambda)$ for small $\lambda \in \Lambda^{reg}$. Then by Corollary 3.2, we obtain the isomorphism.

5.3. **The compactness theorem.**

Definition 5.8. Let $p, q \in \mathrm{Crit}(\mathcal{F})$. Let $\lambda_\nu \in \mathbb{R}^+$ be a sequence such that $\lim_{\nu \to \infty} \lambda_\nu = 0$. We say that a sequence of (parametrized) trajectory $\widetilde{x}_\nu \in \widetilde{\mathcal{M}}^{\lambda_\nu}(p,q)$ *converges* to a parametrized fast-slow solution $\mathscr{X} = (p_0, \sigma_1, p_1, \sigma_2, \ldots, p_{n-1}, \sigma_n, p_n) \in \widetilde{\mathcal{N}}^0(p,q)$, if the following holds

1. For any fast solution σ_i contained in $\mathscr{X}$, there exists $t_{i,\nu} \in \mathbb{R}$ such that the sequence of maps
$$\widetilde{x}_\nu(\cdot + t_{i,\nu}) = (x_\nu(\cdot + t_{i,\nu}), \eta_\nu(\cdot + t_{i,\nu}))$$
converges to σ_i in the C^∞_{loc}-topology.
2. For any slow solution σ_j contained in $\mathscr{X}$, there exists a sequence of intervals $I_{j,\nu} \subset \mathbb{R}$ such that
$$\lim_{\nu \to \infty} d(I_{j,\nu}, t_{i,\nu}) = \infty$$
for any $t_{i,\nu}$ satisfying the first condition for any fast solution σ_i, and such that
$$\lim_{\nu \to \infty} \widetilde{x}_\nu(I_{j,\nu}) = \sigma_j$$
in Hausdorff topology.

It is easy to see that the limit is unique in $\mathcal{N}^0(p,q)$ and it only depends on the sequence of orbits $[\widetilde{x}_\nu]$ but not the representatives. Moreover, the image of $\widetilde{x}_\nu$ will converge to the image of $\mathscr{X}$ in Hausdorff topology.

We would like to prove the following theorem in the remaining of this section.

Theorem 5.9. *Suppose $p, q \in \mathrm{Crit}(\mathcal{F})$ and $\lambda_\nu \to 0^+$ be a sequence of real numbers. Suppose $\widetilde{x}_\nu = (x_\nu, \eta_\nu) \in \widetilde{\mathcal{M}}^{\lambda_\nu}(p,q)$. Then there exists a subsequence (still indexed by ν), and a fast-slow solution $\mathscr{X} = (p_0, \sigma_1, p_1, \sigma_2, \cdots, p_{n-1}, \sigma_n, p_n) \in \widetilde{\mathcal{N}}^0(p,q)$, such that (x_ν, ξ_ν) converges to $\mathscr{X}$ in the sense of Definition 5.8.*

A key point in proving this theorem is to use the energy control to bound the number of pieces of fast trajectories appearing in the limit. This is similar to the proof of Gromov compactness theorem of J-holomorphic curves where there is a lower bound of energy for any nontrivial bubble. After finding all the fast trajectories in the limit, each adjacent two of them are connected by a slow trajectory. And before the first fast trajectory and after the last fast trajectory, one may or may not need to add nontrivial slow ones, depending on the types of p and q (i.e., repeller or attracter of the slow flow).

From now on the sequence $\widetilde{x}_\nu$ is given and we are free to take subsequences as many times as we want.

5.3.1. *The limit set.* For any subsequence ν' of ν and intervals $I_{\nu'} \subset \mathbb{R}$, set

$$L_\infty(\nu', I_{\nu'}) = \bigcap_{k\geq 1} \overline{\cup_{\nu'\geq k} \widetilde{x}_{\nu'}(I_{\nu'})} \subset M \times \mathbb{R}. \tag{5.3}$$

Lemma 5.10. *We have*

(I) *$(x, \eta) \in L_\infty(\nu', I_{\nu'})$ if and only if there exists a subsequence ν'' of ν' and a sequence of numbers $t_{\nu''} \in I_{\nu''}$, such that $\lim_{\nu''\to\infty} \widetilde{x}_{\nu''}(t_{\nu''}) = (x, \eta)$.*

(II) *$L_\infty(\nu', I_{\nu'})$ is compact.*

(III) *If there exist $s_{\nu'} \in I_{\nu'}$ such that $\lim_{\nu'\to\infty} \widetilde{x}_{\nu'}(s_{\nu'})$ exists, then $L_\infty(\nu', I_{\nu'})$ is connected.*

Proof. The first statement and the closedness is by definition, and compactness follows from Lemma 2.3. It remains to show the connectedness. If not, then there are two nonempty closed subset $L_1, L_2 \subset M \times i\mathbb{R}$ such that $L_\infty(\nu', I_{\nu'}) = L_1 \cup L_2$, and L_1, L_2 has a nonzero distance. Suppose $\lim_{\nu'\to\infty} \widetilde{x}_{\nu'}(s_{\nu'}) \in L_1$, then there exists a subsequence (still indexed by ν') and $w_{\nu'} \in I_{\nu'}$ such that $d(\widetilde{x}_{\nu'}(w_{\nu'}), L_2) \to 0$. Then there exists $w'_{\nu'} \in [s_{\nu'}, w_{\nu'}]$ such that $d(\widetilde{x}_{\nu'}(w'_{\nu'}), L_1) = d(\widetilde{x}_{\nu'}(w'_{\nu'}), L_2) > 0$ for the same subsequence. Then there is a subsequence of $\widetilde{x}_{\nu'}(w'_{\nu'})$ converges to a point "between" L_1 and L_2, which contradicts with $L_\infty(\nu', I_{\nu'}) = L_1 \cup L_2$. □

The following lemma is obvious.

Lemma 5.11. *suppose we have two sequence of numbers, $s_\nu < t_\nu$ with $\lim_{\nu\to\infty} \widetilde{x}_\nu(s_\nu) = a$, $\lim_{\nu\to\infty} \widetilde{x}_\nu(t_\nu) = b$. Then $\mathcal{F}(a) \geq \mathcal{F}(b)$. And for any $z \in L_\infty(\nu, [s_\nu, t_\nu])$, $\mathcal{F}(z) \in [\mathcal{F}(a), \mathcal{F}(b)]$.*

5.3.2. *Identify all fast orbits.* By the compactness of $L_\infty(\nu, \mathbb{R})$, and the fact that $\lambda_\nu \to 0$, for any sequence $t_\nu \in \mathbb{R}$, there exists a subsequence for which $\eta_\nu(\cdot + t_\nu)$ converges to a constant function η_1, on any finite interval. Then by usual argument of Morse theory, there is a subsequence of $\widetilde{x}_\nu(\cdot + t_\nu)$ converging to a fast solution $\widetilde{y}_1 := (y_1, \eta_1) : \mathbb{R} \to M \times \mathbb{R}$ with $\mathrm{Im}\widetilde{y}_1 \subset M \times \{\eta_1\}$, in C^∞_{loc}-topology. So $L_\infty(\nu, \mathbb{R})$ is the union of fast orbits (which could be constant orbits). We first prove that for a suitable subsequence it only contains finitely many nonconstant ones.

Proposition 5.12. *There exists a subsequence (still indexed by ν without loss of generality), and $t_{1,\nu}, \ldots, t_{n,\nu} \in \mathbb{R}$ (n could be zero), satisfying*

(I) *$t_{1,\nu} < t_{2,\nu} < \cdots < t_{n,\nu}$, $\lim_{\nu\to\infty} |t_{i,\nu} - t_{j,\nu}| = \infty$ for all $i \neq j$.*

(II) *$\widetilde{x}_\nu(\cdot + t_{i,\nu})$ converges to a (nonconstant) fast solution $\sigma_i := (y_i, \eta_i) : \mathbb{R} \to M \times \mathbb{R}$ in the C^∞_{loc}-topology.*

(III) *For any sequence $s_\nu \in \mathbb{R}$ such that $\lim_{\nu\to\infty} |s_\nu - t_{i,\nu}| = \infty$ for all i, any convergent subsequence of $\widetilde{x}_\nu(s_\nu)$ has limit in $\mathcal{C}_\mathcal{F}$.*

Proof. If $L_\infty(\nu, \mathbb{R}) \subset \mathcal{C}_\mathcal{F}$, then nothing has to be proven. Suppose it is not the case. Then we take $\delta, \epsilon > 0$ as in Lemma 4.3 and small enough, and $r > 0$ small enough such that

- All ϵ-short orbits are contained in $\cup_{w\in\mathcal{C}^{sing}_\mathcal{F}} B_r(w)$;

- All nontrivial fast orbits which are not ϵ-short has energy no less than δ;
- For each $w \in \mathcal{C}_{\mathcal{F}}^{sing}$, $\mathcal{C}_{\mathcal{F}} \cap B_{2r}(w)$ can be parametrized using the canonical parametrization as in (4.10), so that $\mathcal{F}$ restricted to both components of $\mathcal{C}_{\mathcal{F}} \cap B_{2r}(w) \setminus \{w\}$ is monotonic.

Let $N = \lfloor \frac{E}{\delta} \rfloor$, where $E = \mathcal{F}(p_-) - \mathcal{F}(p_+)$ is the total energy. Then it is easy to prove by induction, that there exists $k \leq N$ (could be zero), a subsequence ν' of the original sequence, a sequence of numbers $\{t_{i,\nu'}\}_{i=1,\dots,k}$ and fast solutions $\widetilde{y}_i := (y_i, \eta_i)$ satisfying

1. $\lim_{\nu' \to \infty} t_{i+1,\nu'} - t_{i,\nu'} = +\infty$, for $i = 1, \dots, k-1$;
2. $\lim_{\nu' \to \infty} \widetilde{x}_{\nu'}(t_{i,\nu'} + \cdot) = (y_i, \eta_i)$ in C^∞_{loc}-topology;
3. each $\widetilde{y}_i$ has energy $\geq \delta$.
4. Any other nonconstant fast orbit contained in $L_\infty(\nu', \mathbb{R})$ is an ϵ-short orbit.

We replace ν' by ν for simplicity. Let $N' = \# \mathcal{C}_{\mathcal{F}}^{sing}$. Then we can continue the induction to find all ϵ-short orbits, and we claim that the induction stops at finite time and we can find at most $N'(N+1)$ ϵ-short trajectories. Suppose not, then there exists a $w \in \mathcal{C}_{\mathcal{F}}^{sing}$ and a subsequence, still indexed by ν, and sequence of numbers

$$s_{1,\nu} < s_{2,\nu} < \cdots < s_{N+2,\nu} \tag{5.4}$$

such that for each j, $\widetilde{x}_\nu(s_{j,\nu} + \cdot)$ converges in C^∞_{loc} to an ϵ-short solution $\widetilde{z}_j := (z_j, \xi_j)$ whose orbit is contained in $B_r(w)$. And there exists $j \in \{1, \dots, N+1\}$ such that $L_\infty(\nu, [s_{j,\nu}, s_{j+1,\nu}])$ contains no orbits which are not ϵ-short. Then we take $[t_{j,\nu}, t_{j+1,\nu}] \subset [s_{j,\nu}, s_{j+1,\nu}]$ such that

$$\lim_{\nu\to\infty} \widetilde{x}_\nu(t_{j,\nu}) = \lim_{t\to+\infty} \widetilde{z}_j(t) =: a_+ \in \mathcal{C}_{\mathcal{F}}, \ \lim_{\nu\to\infty} \widetilde{x}_\nu(t_{j+1,\nu}) = \lim_{t\to-\infty} \widetilde{z}_{j+1}(t) =: b_- \in \mathcal{C}_{\mathcal{F}}$$

and denote $\lim_{t\to-\infty} \widetilde{z}_j(t) = a_-$, $\lim_{t\to+\infty} \widetilde{z}_{j+1}(t) = b_+$. Suppose $L_\infty(\nu, [t_{j,\nu}, t_{j+1,\nu}])$ contains some other ϵ-short orbit contained in $B_r(w)$, then it can contain only finitely many, because those orbits must start from the arc between a_- and b_- and ends at the arc between a_+ and b_+, whose energy has a nonzero lower bound. Therefore, by taking a subsequence if necessary, and chooing subintervals of $[t_{j,\nu}, t_{j+1,\nu}]$, we may assume that $T = L_\infty(\nu, [t_{j,\nu}, t_{j+1,\nu}])$ contains no other ϵ-short orbit which is also contained in $B_r(w)$. This implies that

$$T \subset \mathcal{C}_{\mathcal{F}} \cup \bigcup_{v \in \mathcal{C}_{\mathcal{F}}^{sing},\ v \neq w} B_r(v).$$

Indeed, T is contained in a connected component of the latter set, which is homotopy equivalent to a circle or $\mathbb{R}$.

Now by the fact that w is a nondegenerate critical point of $\mathcal{F}|_{\mathcal{C}_{\mathcal{F}}}$, we see that $w \notin T$, because otherwise by Lemma 5.11, $\mathcal{F}(w) \in [\mathcal{F}(a_+), \mathcal{F}(b_-)]$. But there are other points contained in $\mathcal{C}_{\mathcal{F}} \cap (B_{2r}(w) \setminus B_r(w))$ which is not in T, by the monotonicity requirement in Lemma 5.11. Then this contradicts with the connectedness of T.

□

5.3.3. *Identify all slow orbits.* One can not go from p to q only through fast orbits because $\eta_p \neq \eta_q$ according to our assumption (A12). Hence in the limit slow orbits appear. The identification of them is easy intuitively, because $\mathcal{C}_{\mathcal{F}}$ is 1-dimensional and our limit object should be connected.

Proposition 5.13. *Assume the sequence $\widetilde{x}_\nu$ and $\{t_{i,\nu}\}_{i=1,\dots,n}$ satisfy the condition of the last proposition. Let $t_{0,\nu} = -\infty$ and $t_{n+1,\nu} = +\infty$. Then there exists a subsequence (still indexed by ν), and slow orbits $\gamma_i \subset \mathcal{C}_\mathcal{F}$, $i = 0, \dots, n$ satisfying the following:*

- (I) *For each $i = 1, \dots, n$, γ_i is from $\widetilde{y}_i(+\infty)$ to $\widetilde{y}_{i+1}(-\infty)$.*
- (II) *If $p \in \mathrm{Crit}^+(\mathcal{F})$, then γ_0 is the single point p; if $q \in \mathrm{Crit}^-(\mathcal{F})$, then γ_n is the single point q.*
- (III) *For each $i = 1, \dots, n-1$, there exists intervals $(a_{i,\nu}, b_{i,\nu}) \subset (t_{i-1,\nu}, t_{i,\nu})$ such that $\lim_{\nu\to\infty} |a_{i,\nu} - t_{i-1,\nu}| = \lim_{\nu\to\infty} |b_{i,\nu} - a_{i,\nu}| = \lim_{\nu\to\infty} |t_{i,\nu} - b_{i,\nu}| = \infty$ and $\widetilde{x}_\nu([a_{i,\nu}, b_{i,,\nu}])$ converges to γ_i in Hausdorff topology.*

Proof. By the previous proposition, we may find sequences of intervals $I_{i,\nu} = [a_{i,\nu}, b_{i,\nu}] \subset \mathbb{R}$, $i = 1, \dots, n-1$ such that for each ν, $I_{i,\nu} \cap I_{j,\nu} = \emptyset$, for $i \neq j$, $b_{i,\nu} - a_{i,\nu} \to \infty$, and

$$\lim_{\nu\to\infty} \widetilde{x}_\nu(a_{i,\nu}) = \widetilde{y}_i(+\infty), \ \lim_{\nu\to\infty} \widetilde{x}_\nu(b_{i,\nu}) = \widetilde{y}_{i+1}(-\infty).$$

And take $I_{0,\nu} = (-\infty, a_{0,\nu}]$, $I_{n,\nu} = [b_{n,\nu}, +\infty)$ such that

$$\lim_{\nu\to\infty} \widetilde{x}_\nu(a_{0,\nu}) = \widetilde{y}_1(-\infty), \ \lim_{\nu\to\infty} \widetilde{x}_\nu(b_{n,\nu}) = \widetilde{y}_n(+\infty).$$

(1) For each $i \in \{1, \dots, n\}$, set $y_{i,\pm} = \widetilde{y}_i(\pm\infty)$. And set $y_{0,+} = p$, $y_{n+1,-} = q$. For $i = 0, \dots, n$, let $\Gamma_i \subset \mathcal{C}_\mathcal{F}$ be the union of all slow orbits from $y_{i,+}$ to $y_{i+1,-}$. Since $\mathcal{C}_\mathcal{F}$ is a 1-dimensional manifold, Γ_i is either a single point, or is one nontrivial slow orbit, or is the union of two nontrivial slow orbits. We would like to prove that $L_\infty(\nu, I_{i,\nu}) \subset \Gamma_i$.

If Γ_i is the union of two slow orbits, then Γ_i the whole connected component of it in $\mathcal{C}_\mathcal{F}$. By the connectedness of $L_\infty(\nu, I_{i,\nu})$ the claim is true.

Suppose Γ_i is simply-connected and $L_\infty(\nu, I_{i,\nu})$ is also simply-connected. Suppose in this case the claim is not true, then there exists a subsequence (still indexed by ν) and $t_\nu \in I_{i,\nu}$ such that $\lim_{\nu\to\infty} \widetilde{x}_\nu(t_\nu) = z \in L_\infty(\nu, I_{i,\nu}) \setminus \Gamma_i$ and

$$\mathcal{F}(y_{i,+}) > \mathcal{F}(z) > \mathcal{F}(y_{i+1,-}). \tag{5.5}$$

But since $L_\infty(\nu, [t_\nu, b_{i,\nu}])$ and $L_\infty(\nu, [a_{i,\nu}, t_\nu])$ are connected, either the former contains $y_{i,+}$ or the latter contains $y_{i+1,-}$, either of which contradicts Lemma 5.11.

We claim that it is impossible that Γ_i is homeomorphic to a closed interval and $L_\infty(\nu, I_{i,\nu})$ is homeomorphic to a circle $C \subset \mathcal{C}_\mathcal{F}$. Suppose not, then Lemma 5.11 implies that $y_{i,+}$ and $y_{i+1,-}$ must be the absolute maximum and minimum of the function $\mathcal{F}$ restricted to this component C. The critical points of $\mathcal{F}|_C$ can be put in a cyclic order and adjacent to $y_{i,+}$, the two local minimum are $y_{i+1,-}$ and some $z \in C \setminus \Gamma_i$. Taking a subsequence and find $t_\nu \in I_{i,\nu}$ such that $\lim_{\nu\to\infty} \widetilde{x}_\nu(t_\nu) = z$. And consider the interval $J_{i,\nu} = [t_\nu, b_{i,\nu}]$. Then $L_\infty(\nu, J_{i,\nu})$ is homeomorphic to a closed interval and doesn't contain $y_{i,+}$. But this is impossible because then it must contain another local maximum z' between z and $y_{i+1,-}$ to keep the connectedness of $L_\infty(\nu, J_{i,\nu})$, while $\mathcal{F}(z') \notin [\mathcal{F}(z), \mathcal{F}(y_{i+1,0})]$, which contradicts with Lemma 5.11.

(2) We can identify one slow orbit contained in Γ_i. More precisely, we would like to show that, there exists $\gamma_i \subset \Gamma_i$ a slow orbit from $y_{i,+}$ to $y_{i+1,-}$ and a subsequence ν' such

that $L_\infty(\nu', I_{i,\nu'}) = \gamma_i$. Assume it is impossible, then $\Gamma_i = \gamma_i^1 \cup \gamma_i^2$ the union of two different (nonconstant) slow orbits, and we can assume that there exists a subsequence ν' such that for *any* further subsequence $\{\nu''\} \subset \{\nu'\}$, $\gamma_i^1 \subset L_\infty(\nu'', I_{i,\nu''})$.

Now suppose there exists $\widetilde{z} \in L_\infty(\nu', I_{i,\nu'}) \cap (\Gamma_i \setminus \gamma_i^1)$. Then there exists a subsequence ν'', and $t^-_{\nu''}, t_{\nu''}, t^+_{\nu''} \in I_{i,\nu''}$, $t^-_{\nu''} < t_{\nu''} < t^+_{\nu''}$, such that

$$\lim_{\nu''\to\infty} \widetilde{x}_{\nu''}(t_{\nu''}) = \widetilde{z}, \quad \lim_{\nu''\to\infty} \widetilde{x}_{\nu''}(t^\pm_{\nu''}) = \widetilde{z}^\pm \in \mathrm{Int}(\gamma_i^1)$$

and $\mathcal{F}(y_{i,+}) > \mathcal{F}(\widetilde{z}^-) > \mathcal{F}(\widetilde{z}) > \mathcal{F}(\widetilde{z}^+) > \mathcal{F}(y_{i+1,-})$.

Since

$$\left(\Gamma_i \setminus \gamma_i^1\right) \cap \mathcal{F}^{-1}\left(\left[\mathcal{F}(\widetilde{z}^-), \mathcal{F}(\widetilde{z}^+)\right]\right)$$

has a nonzero distance from γ_1, it means

$$L_\infty(\nu'', [t^-_{\nu''}, t^+_{\nu''}])$$

is disconnected, which contradicts with Lemma 5.10. Hence we have $L_\infty(\nu', I_{i,\nu'}) = \gamma_i^1$.

(3) So in summary, there exists a slow orbit γ_i from $y_{i,+}$ to $y_{i+1,-}$, and a subsequence (still indexed by ν), such that $L_\infty(\nu, I_{i,\nu}) = \gamma_i$. By the connectedness of $L_\infty(\nu, I_{i,\nu})$, this implies that $\widetilde{x}_\nu(I_{i,\nu})$ converges to γ_i in Hausdorff topology. (II) of this proposition simply follows from the requirement that $\mathcal{F}$ is decreasing along γ_0 and γ_n, and p is attracting and q is repelling in this case.

□

So we complete the proof of the compactness theorem.

5.4. **Gluing.** In this subsection we prove the following theorem.

Theorem 5.14. *Suppose $p \in \mathrm{Crit}_k(\mathcal{F})$, $q \in \mathrm{Crit}_{k-1}(\mathcal{F})$. Then, there exists $\epsilon_0 > 0$, such that for all $\lambda \in (0, \epsilon_0]$, there exists a bijection*

$$\Phi^\lambda : \mathcal{N}^0(p,q) \to \mathcal{M}^\lambda(p,q) \tag{5.6}$$

such that for any $\mathscr{X} \in \widetilde{\mathcal{N}}^0(p,q)$, there exist representatives $\widetilde{x}^\lambda \in \widetilde{\mathcal{M}}^\lambda(p,q)$ of $\Phi^\lambda([\mathscr{X}])$ such that $\{\widetilde{x}^\lambda\}$ converges to $\mathscr{X}$ in the sense of Definition 5.8.

Proof. Let $p \in \mathrm{Crit}_k(\mathcal{F})$, $q \in \mathrm{Crit}_{k-1}(\mathcal{F})$. Theorem 5.9 implies that for $\lambda > 0$ small, any orbit in $\mathcal{M}^\lambda(p,q)$ is close to a fast-slow orbit in $\mathcal{N}^0(p,q)$. We must show that for any fast-slow orbit from p to q, if $\lambda > 0$ is small, then there exists a unique orbit from p to q that lies near the given fast-slow orbit.

The proof consists of two parts. In the first part, we construct the gluing map Φ^λ, by using the Exchange Lemma (Lemma A.1) and General Exchange Lemma (Lemma A.2). In the second part, we show that Φ^λ is a bijection.

We remark that, there is only C^r-version but no C^∞-version of the Exchange Lemma and General Exchange Lemma (cf. Section A). Hence the unique orbits we constructed in the first part are only of class C^s *a priori*, for some $0 < s < r$. But since all such orbits are solutions to an ODE with smooth coefficients, they are actually smooth.

Now we start to construct the gluing map. We will frequently use Proposition 5.5 without citing it, and we will use the notation $W^u(p,\lambda)$, etc., from Subsection 2.3.

Let $\mathscr{X} = (p_0, \sigma_1, p_1, \dots, p_{n-1}, \sigma_n, p_n)$ be a fast-slow solution from p to q. For small $\lambda > 0$, we will follow $W^u(p,\lambda)$ around the fast-slow solution until it meets $W^s(q,\lambda)$ transversally. Any orbit in $\mathcal{M}^\lambda(p,q)$ that is near the given fast-slow solution must be in the portion of $W^u(p,\lambda)$ that we follow; uniqueness is a consequence of the transversality at the end of the proof (Step 8).

We have $p \in \mathrm{Crit}_k^+(\mathcal{F}) \cup \mathrm{Crit}_k^-(\mathcal{F})$ and $q \in \mathrm{Crit}_{k-1}^+(\mathcal{F}) \cup \mathrm{Crit}_{k-1}^-(\mathcal{F})$.

Step 1. Suppose $p = (x_p, \eta_p) \in \mathrm{Crit}_k^+(\mathcal{F})$. Then $\mathrm{index}\,(x_p, f_{\eta_p}) = k$, and σ_1 is a fast solution from p to $p_1 = (x_{p_1}, \eta_{p_1}) \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, with $\mathrm{index}\,(x_{p_1}, f_{\eta_{p_1}}) = k-1$. By (A13), $W^u(p,0)$ meets $W^s(p_1,0)$ transversally within $M \times \{\eta_p\}$ along σ_1. Then $W^u(p,0)$ meets $W^s(\sigma_2,0)$ transversally in $M \times \mathbb{R}$ along σ_1. (More precisely, σ_2 should be replaced by a compact portion of the complete orbit corresponding to σ_2; we shall use this sort of abuse of notation throughout this section.) The dimension of the intersection is $\dim W^u(p,0) + \dim W^s(\sigma_2,0) - (n+1) = k + (n-(k-1)+1) - (n+1) = 1$.

Thus σ_1 is isolated in the intersection. The next orbit σ_2 is slow, and we have (i) $p_2 \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, or (ii) $p_2 \in \mathcal{C}_\mathcal{F}^{sing}$. In both cases, $\mathrm{index}\,(x_{p_2}, f_{\eta_{p_2}}) = k-1$.

Step 2. After Step 1, in case (i), the Exchange Lemma [9] implies that for $\lambda > 0$ small, $W^u(p,\lambda)$, followed along the flow, becomes close to $W^u(\sigma_2,0)$ near p_2. By *close* we mean close in the C^s-topology for some large s, which decreases in the course of the proof; see Appendix A. Notice that the dimension of $W^u(\sigma_2,0)$ is $(k-1)+1 = k$ as it should be.

The next orbit σ_3 is fast, and we have (i) $p_3 \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, or (ii) $p_3 \in \mathcal{C}_\mathcal{F}^{sing}$. In case (i), $\mathrm{index}\,(x_{p_3}, f_{\eta_{p_3}}) = k-1$; in case (ii), $\mathrm{index}\,(x_{p_3}, f_{\eta_{p_3}}) = k-2$.

Step 3. After Step 1, in case (ii), let N_0 denote the center manifold of (1.6)–(1.7) for $\lambda = 0$ at p_2, which has dimension 2. The choice of center manifold is not unique; we may choose it to include the start of σ_3 (see Appendix B). By the Exchange Lemma, for $\lambda > 0$ small, $W^u(p,\lambda)$, followed along the flow, becomes close to $W^u(\sigma_2,0)$ where σ_2 enters N_0. Of course $W^u(\sigma_2,0)$ (dimension k) is transverse to $W^s(N_0,0)$ (dimension $(n-(k-1)-1)+2 = n-k+2$); $n-(k-1)-1$ is the number of negative eigenvalues at p_2. The dimension of the intersection is 1. For small $\lambda > 0$, N_0 perturbs to a normally hyperbolic invariant manifold N_λ, and $W^u(p,\lambda)$ is transverse to $W^s(N_\lambda,\lambda)$.

The solution in $W^u(p,\lambda) \cap W^s(N_\lambda,\lambda)$ approaches a solution $p_2^\lambda(t)$ in N_λ that is initially near $\sigma_2 \cap N_0$. The system restricted to N_λ, $\lambda \geq 0$, has been analyzed in [12]; see Appendix A. The result is that $p_2^\lambda(t)$ leaves N_λ close to $\sigma_3 \cap N_0$. Then the General Exchange Lemma implies that $W^u(p,\lambda)$ becomes close to the restriction of $W^u(N_0,0)$ to $\sigma_3 \cap N_0$, i.e., $W^u(p_2,0)$, as $W^u(p,\lambda)$ exits a neighborhood of N_λ. $W^u(p_2,0)$ has dimension $(k-1)+1 = k$ as it should. The orbit σ_3 is fast, $p_3 \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, and $\mathrm{index}\,(x_{p_3}, f_{\eta_{p_3}}) = k-1$.

Step 4. After Step 2, in case (i), $W^u(p,\lambda)$ is close to $W^u(\sigma_2,0)$ near p_2, and by (A10), $W^u(\sigma_2,0)$ is transverse to $W^s(\sigma_4,0)$ along σ_3. Continuation from here is like continuation after Step 1, described in Steps 2–3.

Step 5. After Step 2, in case (ii), we note that by (A13), $W^u(p_2,0)$ (dimension $k-1$) is transverse to $W^s(p_3,0)$ (dimension $n-(k-2)$) within $M \times \{\eta_{p_3}\}$ along σ_3. Let N_0 denote the center manifold of (1.6)–(1.7) for $\lambda = 0$ at p_3, which has dimension 2; we choose it to include the end of σ_3 (see Appendix B). Then $W^u(\sigma_2, 0)$ (dimension k) is transverse to $W^s(N_0, 0)$ (dimension $(n-(k-2)-1)+2 = n-k+3$) along σ_3. The intersection is 2-dimensional and consists of solutions that track an open set of solutions in N_0 around σ_3.

Step 6. After Step 3, we note that by (A13), $W^u(p_2, 0)$ (dimension k) meets $W^s(p_3, 0)$ (dimension $n-(k-1)$) transversally within $M \times \{\eta_{p_3}\}$ along σ_3. As in Step 1, $W^u(p_2, 0)$ meets $W^s(\sigma_4, 0)$ transversally in $M \times \mathbb{R}$ along σ_3, and $W^u(p, \lambda)$ is close to $W^u(p_2, 0)$ near σ_3. The next orbit σ_4 is a slow solution from p_3 to (i) $p_4 \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, or (ii) $p_4 \in \mathcal{C}_\mathcal{F}^{sing}$. In both cases, $\operatorname{index}(x_{p_4}, f_{\eta_{p_4}}) = k-1$. Continuation from here is like continuation after Step 1, described in Steps 2–3.

Step 7. After Step 5, for small $\lambda > 0$, the tracked orbits include an open set U_λ in N_λ, the perturbation of N_0, that lies above an open set U_0 in N_0 that contains $\sigma_4 \cap N_0$ in its interior. The General Exchange Lemma implies that $W^u(p, \lambda)$ is close to the restriction of $W^u(N_0, 0)$ to U_0, i.e., to $W^u(\sigma_4 \cap N_0)$, as it exits a neighborhood of N_λ. Continuation from here is like continuation after Step 2, described in Steps 4–5.

Step 8. Continuation proceeds using analogs of the steps previously described until $W^u(p)$ arrives near p_{n-1}.

If $q \in \mathrm{Crit}_k^+(\mathcal{F})$, then σ_n is slow, and $\operatorname{index}(x_{p_{n-1}}, f_{\eta_{p_{n-1}}}) = \operatorname{index}(x_q, f_{\eta_q}) = k-1$. $W^u(p, \lambda)$ arrives near p_n along the slow orbit σ_n, close to $W^u(\sigma_n, 0)$, which has dimension k. $W^u(\sigma_n, 0)$ is transverse to $W^s(\sigma_n, 0)$, with dimension $n-(k-1)+1 = n-k+2$. Therefore $W^u(p, \lambda)$ is transverse to $W^s(q, \lambda)$, which is close to $W^s(\sigma_n, 0)$. The intersection has dimension 1 and gives the solution from p to q.

If $q \in \mathrm{Crit}_k^-(\mathcal{F})$, then σ_n is a fast orbit from $p_{n-1} \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$ to $q \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$; $\operatorname{index}(x_{p_{n-1}}, f_{\eta_{p_{n-1}}}) = k-1$ and $\operatorname{index}(x_q, f_{\eta_q}) = k-2$. By (A11), $W^u(p_{n-1}, 0)$ meets $W^s(q, 0)$ transversally in $M \times \{\eta_q\}$ along σ_n. Therefore $W^u(\sigma_n, 0)$ meets $W^s(q, 0)$ transversally in $M \times \mathbb{R}$ along σ_n. Since $W^u(p, \lambda)$ is close to $W^u(\sigma_n, 0)$, $W^u(p, \lambda)$ meets $W^s(q, \lambda)$ transversally in $M \times \mathbb{R}$ near σ_n. The intersection has dimension one and gives the solution from p to q.

Step 9. Suppose $p = (x_p, \eta_p) \in \mathrm{Crit}_k^-(\mathcal{F})$. Then $\operatorname{index}(x_p, f_{\eta_p}) = k-1$, and σ_1 is a slow solution from p to $p_1 = (x_{p_1}, \eta_{p_1})$. Either (i) $p_1 \in \mathcal{C}_\mathcal{F} \setminus \mathcal{C}_\mathcal{F}^{sing}$, or (ii) $p_1 \in \mathcal{C}_\mathcal{F}^{sing}$. In either case, $\operatorname{index}(x_{p_1}, f_{\eta_{p_1}}) = k-1$.

In case (i), $W^u(p, \lambda)$ is close to $W^u(\sigma_1, 0)$. Step 3 above describes how to continue from p_1. Case (ii) is left to the reader.

Now we show that the gluing map just constructed is a bijection. We first note that there exist small $\delta > 0$ and $\lambda_0 > 0$ such that for all $\lambda \in (0, \lambda_0)$, the orbit we constructed is the unique one in $\mathcal{M}^\lambda(p, q)$ such that its Hausdorff distance from the fast-slow orbit $[\mathscr{X}]$ is less than δ. This is a general fact about exchange lemma constructions; see for example [10], p. 1021.

Thus the gluing map is a bijection provided it is surjective. We claim that there exists $\lambda_0 > 0$ such that the gluing map is surjective for all $\lambda \in (0, \lambda_0)$. If not, then there exist a sequence $\lambda_i \to 0$ and a sequence of orbits $[\widetilde{x}_i] \in \mathcal{M}^{\lambda_i}(p, q)$ such that for each i, $[\widetilde{x}_i]$ is not in the image of Φ^{λ_i}. Then, by the compactness theorem (Theorem 5.9), without loss of generality, $[\widetilde{x}_i]$ converges to a fast-slow orbit $[\mathscr{Y}]$ in the sense of Definition 5.8. In particular, this sequence converges in the Hausdorff topology. Then for large i, $\Phi^{\lambda_i}([\mathscr{Y}])$, which is the unique orbit in $\mathcal{M}^{\lambda_i}(p,q)$ in a small Hausdorff neighborhood of $[\mathscr{Y}]$, must be $[\widetilde{x}_i]$. This contradiction proves the claim. □

6. Basic Morse theory and the chain complex $\mathcal{C}^0$

Let us recall the following basic properties of any Morse function $\mu : M \to \mathbb{R}$, where M is a compact manifold.

(1) If $[c_1, c_2]$ contains no critical value of μ, then $\mu^{-1}((-\infty, c_1])$ is diffeomorphic to $\mu^{-1}((-\infty, c_2])$;

(2) If (i) $c \in (c_1, c_2)$ is the only critical value of μ contained in $[c_1, c_2]$, (ii) there is a unique critical point x of μ such that $\mu(x) = c$, and (iii) x has Morse index k, then the sublevel set $\mu^{-1}((-\infty, c_2])$ is obtained from the sublevel set $\mu^{-1}((-\infty, c_1])$ by attaching a k-handle.

Thus a change in the level set of μ corresponds to a change in the sublevel set of μ. The chain complex $\mathcal{C}^0$ defined by counting fast-slow orbits gives an indirect way to obtain, at the homology level, this change of topology. In this section we will give a brief indication of how this works.

We replace μ by $\mu_c = \mu - c$ for any regular value c of μ, and denote the corresponding Lagrange multiplier by $\mathcal{F}_c : M \times \mathbb{R} \to \mathbb{R}$. Then $\mathrm{Crit}(\mathcal{F}_c)$ is just the critical point set of the restriction of f to the hypersurface $\mu^{-1}(c)$. On the other hand, the slow manifold $\mathcal{C}_{\mathcal{F}_c} = \mathcal{C}_{\mathcal{F}}$ is unchanged, and $\mathrm{Crit}(\mathcal{F}_c) = \mathcal{C}_{\mathcal{F}_c} \cap (\mu^{-1}(c) \times \mathbb{R})$. The fast flow is unchanged when we vary c, and the changes in generators and the slow flow can be seen transparently from the intersection of $\mathcal{C}_{\mathcal{F}}$ and $\mu^{-1}(c) \times \mathbb{R}$.

We have to consider various "wall-crossing" phenomena of the chain complex $\mathcal{C}^0(f, \mu - c)$ when we vary c, not only at the critical values of μ, but also at each c where an assumption we made to define the chain complex fails. We assume that for each $c \in \mathbb{R}$, if c is a critical value of μ, then $\mu^{-1}(c)$ contains only one critical point; if c is a regular value, then at most one of our assumptions is violated; and the violations happen at discrete values of c. This assumption is satisfied for generic (f, μ, g).

6.1. When c does not cross a critical value.

When $[c_1, c_2]$ contains no critical value of μ, $\mu^{-1}(c_1)$ and $\mu^{-1}(c_2)$ are diffeomorphic. We look at the chain complexes $\mathcal{C}^0(f, \mu - c_1)$ and $\mathcal{C}^0(f, \mu - c_2)$ in several cases.

1. Suppose that as c moves from c_1 to c_2, a degenerate critical point of $\mu|_{\mathcal{C}_{\mathcal{F}}}$ appears (in violation of Proposition 4.4) and then splits into nondegenerate critical points. Generically two nondegenerate critical points are produced as shown in Figure 3. Thus two new generators of $\mathcal{C}^0(f, \mu - c_2)$ are produced. We denote them by $p \in \mathrm{Crit}_k^+$ and $q \in \mathrm{Crit}_{k+1}^-$. Between them there is a slow trajectory γ. It is easy to check that the two chain complexes are quasi-isomorphic.



Figure 3. The dashed lines represent the hypersurfaces $\mu = c_1$ and $\mu = c_2$, one of which has two more intersection points with $\mathcal{C}_{\mathcal{F}}$ than the other. This change introduces two new generators in the chain complex, but they do not affect its homology.

2. Suppose that as c moves from c_1 to c_2, a critical point of $\mathcal{F}_c$ passes through the start or end of a handle-slide (in violation of (A5)). See Figure 4. It is easy to see that $\mathcal{C}^0(f, \mu - c_1)$ and $\mathcal{C}^0(f, \mu - c_2)$ are actually isomorphic.



Figure 4. The dashed lines represent the hypersurfaces $\mu = c_1$ and $\mu = c_2$. σ is a handle-slide. This change replace a generator $p \in \mathrm{Crit}_k^-$ by another $q \in \mathrm{Crit}_k^-$ and actually gives isomorphic chain complexes.

3. Suppose that as c moves from c_1 to c_2, a critical point $p \in \mathrm{Crit}_k^-$ passes through a point $w \in \mathcal{C}_{\mathcal{F}}^{sing}$ (in violation of (A5)) and then becomes a critical point $q \in \mathrm{Crit}_k^+$; see Figure 5. If there is a cusp trajectory σ_1 starting at w, it will contribute to the definition of $\partial^0(p)$ in the chain complex $\mathcal{C}^0(f, \mu - c_1)$. In $\mathcal{C}^0(f, \mu - c_2)$, the corresponding contribution to $\partial^0(q)$ is given by a fast trajectory σ_2 starting from q.

6.2. **When c crosses a critical value.** Suppose that one point $c \in (c_1, c_2)$ is a critical value μ, with a single critical point x of index k of μ. Then topologically, $\mu^{-1}(c_1)$ and $\mu^{-1}(c_2)$ differ by attaching a k-handle, and the homology changes accordingly. We can recover this fact by comparing the chain complexes $\mathcal{C}^0(f, \mu - c_1)$ and $\mathcal{C}^0(f, \mu - c_2)$. In the generic case, one complex has two more generators, one of index $k + 1$ and one of index $n - k$. Figure 6 illustrates one case of this changing.



FIGURE 5. The dashed lines sides represent the hypersurfaces $\mu = c_1$ and $\mu = c_2$ respectively. σ_1 is a cusp trajectory starting from $w \in \mathcal{C}_{\mathcal{F}}^{sing}$, and σ_2 is a regular fast trajectory starting from q. The generator $p \in \text{Crit}_k^-$ is replaced by $q \in \text{Crit}_k^+$; the chain complexes are isomorphic.



FIGURE 6. On the left: the left branch of $\mathcal{C}_{\mathcal{F}}$ has index $n-k$ (of the fast flow) and $\eta << 0$, and the right branch of $\mathcal{C}_{\mathcal{F}}$ has index k and $\eta >> 0$. It has two more generators p and q with degrees $n-k$ and $k+1$ respectively than the right picture.

APPENDIX A. GEOMETRIC SINGULAR PERTURBATION THEORY

Readers familiar with geometric singular perturbation theory may wish to skip most of this appendix; however, the paragraph before Theorem A.2 and that theorem may be unfamiliar, and the extension of that theorem described toward the end of this section, while just an observation, is new and essential to this paper.

Let Z be a manifold, let $\dot{p} = f(p, \epsilon) = f_\epsilon(p)$ be a differential equation on Z with parameter ϵ (i.e., each f_ϵ is a section of TZ), let ϕ_t^ϵ be the flow, and let N_0 be a compact submanifold of Z that is invariant under the the flow of $\dot{p} = f_0(p)$. N_0 is called *normally hyperbolic* if there is a splitting of $TZ|N_0$, $TZ|N_0 = S \oplus U \oplus TN_0$, such that under $D\phi^0_t$, all vectors in S shrink at a faster exponential rate than any vector in TN_0, and under $D\phi_{-t}$, all vectors in U shrink at a faster exponential rate than any vector in TN_0. (There are less restrictive definitions, but this one suffices for our purposes.)

Normally hyperbolic invariant manifolds have stable and unstable manifolds with flow-preserved fibrations, and the whole structure persists under perturbation. This structure is most easily described in local coordinates.

Let us assume that $\dim N_0 = m$, $\dim Z = n + m$, and fibers of S (respectively U) have dimension k (respectively l), with $k + l = n$. Near a point of N_0 one can choose coordinates $p = \Phi(x, y, z, \epsilon)$, $(x, y, z, \epsilon) \in \Omega_1 \times \Omega_2 \times (-\epsilon_0, \epsilon_0)$, Ω_1 open subset of $\mathbb{R}^k \times \mathbb{R}^l$, Ω_2 an open subset of $\mathbb{R}^m$, such that, for small ϵ, $\dot{p} = f(p, \epsilon)$ becomes

$$\dot{x} = A(x, y, z, \epsilon)x, \tag{A.1}$$

$$\dot{y} = B(x, y, z, \epsilon)y, \tag{A.2}$$

$$\dot{z} = h(z, \epsilon) + x^T C(x, y, z, \epsilon)y; \tag{A.3}$$

the matrices A, B, and C are $k \times k$, $l \times l$, and $k \times l$ respectively. One cannot assume that $A(0, 0, z, 0)$ has eigenvalues with negative real part or that $B(0, 0, z, 0)$ has eigenvalues with positive real part, since the exponential convergence of vectors under the linearized flow need only occur as $t \to \pm\infty$. Nevertheless, this is true in the examples below.

We list some facts and terminology.

(1) If $\dot{p} = f(p, \epsilon)$ is C^{r+3}, the coordinate change can be chosen so that the new system is C^{r+1}.

(2) For each ϵ, the subspaces $y = 0$, $x = 0$, and their intersection are invariant. For fixed ϵ, the set $\{(x, y, z) \mid x = 0 \text{ and } y = 0\}$ (dimension m) corresponds to part of a normally hyperbolic invariant manifold N_ϵ; the set $y = 0$ (dimension $m + k$) corresponds to part of the stable manifold of N_ϵ, $W^s(N_\epsilon)$; and the set $x = 0$ (dimension $m + l$) corresponds to part of the unstable manifold of N_ϵ, $W^u(N_\epsilon)$.

(3) If $(x(t), 0, z(t))$ is a solution in $W^s(N_\epsilon)$, then $(0, 0, z(t))$ is a solution in N_ϵ; and if $(0, y(t), z(t))$ is a solution in $W^u(N_\epsilon)$, then $(0, 0, z(t))$ is again a solution in N_ϵ. Each solution in $W^s(N_\epsilon)$ (respectively $W^u(N_\epsilon)$) approaches exponentially a solution in N_ϵ as time increases (respectively decreases).

(4) Given a point $p = (0, 0, z_0)$ in N_ϵ, the *stable* (respectively *unstable*) *fiber* of p is the set of all points $(x, 0, z_0)$ (dimension k) (respectively $(0, y, z_0)$ (dimension l)). For each t, the time-t map of the flow takes fibers to fibers; in this sense the fibration is flow-invariant. Solutions that start in the stable (respectively) unstable fiber of p approach the solution that starts at p exponentially at t increases (respectively decreases).

(5) Given an $P_\epsilon \subset N_\epsilon$, we shall refer to the union of the stable (respectively unstable) fibers of points in P_ϵ as $W^s(N_\epsilon)$ (respectively $W^u(N_\epsilon)$) *restricted to* P_ϵ. If P_ϵ is invariant, $W^s(N_\epsilon)$ restricted to P_ϵ, for example, may be smaller than $W^s(P_\epsilon)$, since the latter may include solutions in N_ϵ that approach P_ϵ at a slower exponential rate, together with points in their stable fibers.

Examples:

(1) Suppose N_0 is a compact manifold of equilibria of dimension m, and each equilibrium in N_0 has k eigenvalues with negative real part and l eigenvalues with positive real part. Then N_0 is a compact normally hyperbolic invariant manifold. The stable (respectively unstable) fiber of a point p in N_0 is just its stable (respectively unstable) manifold. In (A.3), $h(z, 0) \equiv 0$.

(2) Suppose N_0 is a compact manifold with boundary of equilibria of dimension m, and each equilibrium in N_0 has k eigenvalues with negative real part and l eigenvalues with positive real part. N_0 does not fit the definition of a normally hyperbolic invariant manifold that we have given, but a coordinate system as above still exists, except that Ω_2 may have boundary and N_ϵ may be only locally invariant (i.e., solutions may leave through the boundary).

(3) Suppose p_0 is a critical point of $\dot{p} = f_0(p)$ at which the linearization has k eigenvalues with negative real part, l eigenvalues with positive real part, and m eigenvalues with 0 real part. On a neighborhood of p_0, one can choose coordinates as above such that the system takes the form (A.1)–(A.3); this is the Center Manifold Theorem. Here the N_ϵ are locally invariant open manifolds.

(4) Fast-slow systems. Consider the system

$$\dot{w} = f(w, z, \epsilon), \tag{A.4}$$

$$\dot{z} = \epsilon g(w, z, \epsilon), \tag{A.5}$$

with $w \in \mathbb{R}^n$, $z \in R^m$, and $0 \le \epsilon < \epsilon_0$. System (A.4)–(A.5) is a *fast-slow system*; w is the fast variable and z is the slow variable. The fast limit is (A.4)–(A.5) with $\epsilon = 0$. If 0 is a regular value of f, then $\{(w, z) : f(w, z, 0) = 0\}$ is a manifold of dimension m called the *slow manifold*. The slow manifold is the set of equilibria of the fast system. Let N_0 be a compact subset of the slow manifold that is a manifold with boundary of dimension m. N_0 is normally hyperbolic (in the sense of Example 2) if there are numbers k and l with $k + l = n$ such that at each point $(w, z, 0)$ of N_0, $D_z f(w, z, 0)$ has k eigenvalues with negative real part and l eigenvalues with positive real part. By the Implicit Function Theorem, N_0 can be described as $w = \chi(z, 0)$. Then for $\epsilon > 0$ small there is a normally hyperbolic locally invariant manifold N_ϵ (as in Example 2) near N_0, given by $w = \chi(z, \epsilon)$.

The name "geometric singular perturbation theory" comes from the fact that after a rescaling of time, (A.4)–(A.5) can be rewritten as

$$\epsilon \dot{w} = f(w, z, \epsilon), \tag{A.6}$$

$$\dot{z} = g(w, z, \epsilon), \tag{A.7}$$

which is a singularly perturbed system.

In Example 4, N_ϵ is given by an expansion $w = \chi_0(z) + \epsilon \chi_1(z) + \dots$, and the system restricted to N_ϵ, with coordinate z, is given by

$$\dot{z} = \epsilon g(\chi_0(z) + \epsilon \chi_1(z) + \dots, z, \epsilon) = \epsilon g(\chi_0(z), z, 0) + \mathcal{O}(\epsilon^2). \tag{A.8}$$

Thus, after division by ϵ, system (A.4)–(A.5) restricted to N_ϵ is given by $\dot{z} = g(\chi_0(z), z, 0) + \mathcal{O}(\epsilon)$. The differential equation $\dot{z} = g(\chi_0(z), z, 0)$ is called the *slow equation*. The expansion of N_ϵ, and hence of (A.4)–(A.5) restricted to N_ϵ, can be computed to any order using the invariance of N_ϵ. Similar calculations can be done in Examples 2 and 3. In the latter case, typically the center manifold at p_0 for $\epsilon = 0$ is not known explicitly, so it too must be computed as an expansion in a coordinate system near p_0.

We shall abuse terminology and refer to the N_ϵ as normally hyperbolic invariant manifolds in all the above situations.

Suppose we wish to follow an $(l+1)$-dimensional manifold of solutions M_ϵ as it passes near a normally hyperbolic invariant manifold N_ϵ in a manifold Z, where N_0 is a normally hyperbolic invariant manifold of equilibria. We choose coordinates so that the system is (A.1)–(A.3), with $h(z,\epsilon) = \epsilon\tilde{h}(z,\epsilon)$. We assume that near the point $(x^*, 0, z^*)$, the M_ϵ, $\epsilon \geq 0$, fit together to form a C^{r+1} manifold with boundary in $Z \times \mathbb{R}$; and we assume that M_0 meets $W^s(N_0)$ transversally in the solution through $(x^*, 0, z^*)$, which approaches the equilibrium $(0,0,z^*)$ as $t \to \infty$. The system restricted to N_ϵ has the form $\dot{z} = \epsilon\tilde{h}(z,\epsilon)$; we assume $\tilde{h}(z^*,0) \neq 0$. Let ψ_t denote the flow of $\dot{z} = \tilde{h}(z,0)$, and let $z^\dagger = \psi_T(z^*)$ for some $T > 0$. Let $I = \{(\psi_t(z^*) : T - \delta < t < T + \delta\}$, a small interval around $z^\dagger$ in the orbit of z^* for $\dot{z} = \tilde{h}(z,0)$. Let $J = \{0\} \times \{0\} \times I$. Let V be a small open set around $(0,0,z^\dagger)$ in $W^u(N_0)$ restricted to J, which has dimension $l+1$. Then we have:

Theorem A.1 (Exchange Lemma). *For small $\epsilon > 0$, parts of M_ϵ fit together with V to form a C^r manifold in $Z \times \mathbb{R}$.*



(a) (b)

FIGURE 7. The Exchange Lemma with $k = l = m = 1$: (a) $\epsilon = 0$, (b) $\epsilon > 0$.

See Figure 7. Notice that for small $\epsilon > 0$, M_ϵ meets $W^s(N_\epsilon)$ transversally in a solution that tracks a solution in N_ϵ that starts at a point $(0,0,z(\epsilon))$, with $z(\epsilon)$ near z^*. For $\epsilon > 0$, the orbit of $\dot{z} = \epsilon\tilde{h}(z_\epsilon)$ through z_ϵ equals the orbit of $\dot{z} = \tilde{h}(z,\epsilon)$ through $z(\epsilon)$. This orbit is close to the orbit of $\dot{z} = \tilde{h}(z,0)$ through z^*. Thus we find that portions of the orbits tracked in N_ϵ for $\epsilon > 0$ limit on a curve J in N_0 that is not an orbit of the system (A.1)–(A.3) on

N_0 (these orbits are just points). Nevertheless the M_ϵ for $\epsilon > 0$ become close to $W^u(N_0)$ restricted to J.

Now suppose we wish to track an $(l+s+1)$-dimensional manifold of solutions M_ϵ, with $0 \le s \le m-1$, as it passes near a normally hyperbolic invariant manifold N_ϵ in a manifold Z, where N_0 is a normally hyperbolic invariant manifold but does not consist of equilibria. (This occurs, for example, Step 3 of the proof of Theorem 5.14.) We again choose coordinates so that the system is (A.1)–(A.3). We choose a cross-section $\tilde{M}_\epsilon$ to the flow within M_ϵ, of dimension $l+s$ (so that M_ϵ is the union of orbits that start in $\tilde{M}_\epsilon$), and we assume that $\tilde{M}_0$ meets $W^s(N_0)$ transversally at $(x^*, 0, z^*)$. For small $\epsilon \ge 0$, $\tilde{M}_\epsilon$ meets $W^s(N_\epsilon)$ in a manifold Q_ϵ of dimension s; we assume that Q_ϵ projects regularly along the stable fibration to a submanifold P_ϵ of N_ϵ of dimension s. We also assume that for small $\epsilon \ge 0$, the vector field $(0, 0, h(x, \epsilon))$ is not tangent to P_ϵ. (Of course these assumptions follow from the corresponding ones at $\epsilon = 0$.) Finally, we assume that for $\epsilon > 0$, following P_ϵ along the flow for time $\mathcal{O}(\frac{1}{\epsilon})$ produces a submanifold P_ϵ^* of dimension $s+1$ of N_ϵ, and the manifolds P_ϵ^* fit together with a submanifold P_0^* of N_0, of dimension $s+1$, to form a C^{r+1} manifold in $Z \times \mathbb{R}$. This assumption typically requires that some solution of the system restricted to N_0 that starts in P_0 approaches an equilibrium. We emphasize that, analogous to the usual Exchange Lemma, P_0^* is not the result of following P_0 along the flow for $\epsilon = 0$. Let V be a small open neighborhood of P_0^* in $W^u(N_0)$ restricted to P_0^*, which has dimension $l+s+1$.

Theorem A.2 (General Exchange Lemma)**.** *For small $\epsilon > 0$, parts of M_ϵ fit together with V to form a C^r manifold in $Z \times \mathbb{R}$.*

Some technical hypotheses that are not relevant to the present paper have been omitted. We actually need a small generalization of Theorems A.1 and A.2, whose proofs are essentially the same.

We will use C^r-topology to measure submanifolds. If $S \subset X$ is a smooth submanifold, we say another C^r-submanifold S'(of the same dimension) is C^r-close to S, if with some smooth identification of the normal bundle of S with a tubular neighborhood of S, S' can be identified with a C^r-small section of the normal bundle.

In Theorem A.1, replace the assumption that near the point $(x^*, 0, z^*)$, the M_ϵ, $\epsilon \ge 0$, fit together to form a C^{r+1} manifold with boundary, with the assumption that $M_\epsilon \to M_0$ in the C^{r+1}-topology. The conclusion becomes that parts of M_ϵ converge to V in the C^r-topology.

In Theorem A.2, make the replacement in the assumptions just mentioned, and replace the assumption that the P_ϵ^*, $\epsilon \ge 0$, fit together to form a C^{r+1} manifold with the assumption that $P_\epsilon^* \to P_0^*$ in the C^{r+1}-topology. The conclusion again becomes that parts of M_ϵ converge to V in the C^r-topology.

These generalizations are required for the following reason. Consider the fast-slow system

$$\dot{z}_1 = -z_2 + z_1^2, \tag{A.9}$$

$$\dot{z}_2 = -\epsilon, \tag{A.10}$$

which is related to (4.17)–(4.18). See Figure 8. For $\epsilon = 0$, any compact portion N_0 of the curve of equilibria $z_1 = -z_2^{\frac{1}{2}}$ is normally hyperbolic (in fact attracting). It perturbs to the normally hyperbolic manifold N_ϵ, on which system reduces to $\dot{y} = -\epsilon$. For small $\epsilon > 0$, a solution in or close to N_ϵ arrives in the region $\delta < z_1 < 2\delta$ ($\delta > 0$) along a curve given by $z_2 = \rho(z_1, \epsilon)$, $\delta < z_1 < 2\delta$; $\rho = \mathcal{O}(\epsilon^{\frac{2}{3}})$. As $\epsilon \to 0$, this curve, which in examples with greater dimension may be P^*_ϵ, approaches $z_2 = 0$, which in examples may be P^*_0, in the C^s-topology for any s, but does not fit together with $z_2 = 0$ to form a manifold with a high degree of differentiability in $z_1z_2\epsilon$-space.



(a) (b)

Figure 8. Flow of (A.9)–(A.10): (a) $\epsilon = 0$, (b) $\epsilon > 0$.

Appendix B. Choosing the center manifold

Let us consider for concreteness a system in the form (A.1)–(A.3), with N_0 two-dimensional, and the equation on N_ϵ given by (A.9)–(A.10):

$$\dot{x} = A(x, y, z_1, z_2, \epsilon)x, \tag{B.1}$$

$$\dot{y} = B(x, y, z_1, z_2, \epsilon)y, \tag{B.2}$$

$$\dot{z}_1 = -z_2 + z_1^2 + x^T c_1(x, y, z_1, z_2, \epsilon)y, \tag{B.3}$$

$$\dot{z}_2 = -\epsilon + x^T c_2(x, y, z_1, z_2, \epsilon)y, \tag{B.4}$$

$(x, y) \in \mathbb{R}^k \times \mathbb{R}^l$. The system arises by center manifold reduction at the origin, so $A(0, 0, z_1, z_2, 0)$ has eigenvalues with negative real part ,and $B(0, 0, z_1, z_2, 0)$ has eigenvalues with positive real part. We wish to follow a manifold M_ϵ of dimension $l+1$ as it passes N_ϵ (i.e., z_1z_2-space). M_0 meets $W^s(N_0)$ transversally at a point $(x^*, 0.z_1^*, 0)$ with $z_1^* < 0$; hence the intersection includes the semiorbit γ that starts at $(x^*, 0, z_1^*, 0)$, which approaches the origin as $t \to \infty$. We wish to replace the center manifold N_0 for $\epsilon = 0$ by one that contains γ. To do this, replace all semiorbits in Figure 8 that start at points $(0, 0, z_1^*, z_2)$, $|z_2| < (z_1^*)^2$, with the semiorbits that start at $(x^*, 0, z_1^*, z_2)$. (The semiorbits that start at $(0, 0, z_1^*, z_2)$ and at $(x^*, 0, z_1^*, z_2)$ both approach $(0, 0, -z_2^{\frac{1}{2}}, z_2)$ as $t \to \infty$, and the second arrives tangent to z_1z_2-space.) The result will be a new center manifold, in a smaller neighborhood of the origin, that contains γ. The differentiability class of this manifold will decrease as we move away from the origin. Now N_0 perturbs to new center manifold N_ϵ for $\epsilon > 0$.

References

[1] Augustin Banyaga and David Hurtubise, *Morse-Bott homology*, Transactions of the American Mathematical Society **362** (2010), no. 8, 3997–4043.

[2] ———, *Cascades and perturbed Morse-Bott functions*, arXiv: 1110.4609, 2011.

[3] Kai Cieliebak, Ana Gaio, and Dietmar Salamon, *J-holomorphic curves, moment maps, and invariants of Hamiltonian group actions*, International Mathematical Research Notices **16** (2000), 831–882.

[4] Andreas Floer, *Morse theory for Lagrangian intersections*, Journal of Differential Geometry **28** (1988), 513–547.

[5] ———, *The unregularized gradient flow of the symplectic action*, Communications on Pure and Applied Mathematics **41** (1988), no. 6, 775–813.

[6] Urs Frauenfelder, *Floer homology of symplectic quotients and the Arnold-Givental conjecture*, Ph.D. thesis, Swiss federal institute of technology, 2003, pp. 2179–2269.

[7] Ana Gaio and Dietmar Salamon, *Gromov-Witten invariants of symplectic quotients and adiabatic limits*, Journal of Symplectic Geometry **3** (2005), no. 1, 55–159.

[8] Eduardo González and Chris Woodward, *Gauged Gromov-Witten invariants for small spheres*, Mathematische Zeitschrift (To appear).

[9] Christopher Jones, *Geometric singular perturbation theory*, Dynamical systems (R. Johnson, ed.), Lecture Notes in Mathematics, vol. 1609, Springer, 1995.

[10] Christopher K. R. T Jones and Siu-Kei Tin, *Generalized exchange lemmas and orbits heteroclinic to invariant manifolds*, Discrete Contin. Dyn. Syst. Ser. S **2** (2009), no. 4, 967–1023.

[11] Jungsoo Kang, *Invariance property of Morse homology on noncompact manifolds*, arXiv:math/1012.5571.

[12] M. Krupa and P. Szmolyan, *Extending geometric singular perturbation theory to nonhyperbolic points-fold and canard points in two dimensions*, SIAM Journal of Mathematical Analysis **33** (2001), no. 2, 286–314.

[13] Dusa McDuff and Dietmar Salamon, *J-holomorphic curves and symplectic topology*, Colloquium publications, vol. 52, American Mathematical Society, 2004.

[14] John Milnor, *Morse theory*, Annals of Mathematics Studies, no. 51, Princeton University Press, 1963.

[15] Stephen Schecter, *Exchange lemmas. II. General exchange lemma*, Journal of Differential Equations **245** (2008), no. 2, 411–441.

[16] Matthias Schwarz, *Morse homology*, Progress in Mathematics, Birkhauser, 1993.

[17] Edward Witten, *Supersymmetry and Morse theory*, Journal of Differential Geometry **17** (1982), no. 4, 661–692.

[18] ———, *The Verlinde algebra and the cohomology of the Grassmannian*, Geometry, Topology and Physics, International Press, Cambridge, MA, 357-422.

Department of Mathematics, North Carolina State University, Box 8205, Raleigh, NC 27695 USA, 919-515-6533

E-mail address: `schecter@ncsu.edu`

Department of Mathematics, Princeton University, Fine Hall, Washington Road, Princeton, NJ 08544 USA, 312-646-9515

E-mail address: `guangbox@math.princeton.edu`